

Date : June 19, 2009

**$(\ell, 0)$ -JM PARTITIONS AND A LADDER BASED MODEL FOR
THE BASIC CRYSTAL OF $\widehat{\mathfrak{sl}}_\ell$**

CHRIS BERG

ABSTRACT. In [1], Vazirani and I gave a new interpretation of what we called ℓ -partitions, also known as $(\ell, 0)$ -Carter partitions. The primary interpretation of such a partition λ is that it corresponds to a Specht module S^λ which remains irreducible over the finite Hecke algebra $H_n(q)$ when we specialize q to a primitive ℓ^{th} root of unity. In [1], we relied heavily on the description of such a partition in terms of its hook lengths, a condition provided by James and Mathas in [11]. In this paper, I introduce a new description of the crystal $B(\Lambda_0)$ which helps us extend the results of [1] to all $(\ell, 0)$ -JM partitions (similar to $(\ell, 0)$ -Carter partitions, but not necessarily ℓ -regular), by using an analogous condition for hook lengths conjectured by James and Mathas in [17], and recently proven by Fayers in [4]. Using the new description of the crystal, I extend our previous theorems from [1] about the crystal $B(\Lambda_0)$ of $\widehat{\mathfrak{sl}}_\ell$ to a larger class of partitions.

1. INTRODUCTION

The main goal of this paper is to generalize results of [1] to a larger class of partitions. As most of the proofs in the first few sections are generalizations, reading [1] is recommended. One interpretation of the crystal $B(\Lambda_0)$ of $\widehat{\mathfrak{sl}}_\ell$ has as nodes ℓ -regular partitions. In [1] we proved results about where on the crystal $B(\Lambda_0)$ a so-called ℓ -partition could occur. ℓ -partitions are the ℓ -regular partitions for which the Specht modules S^λ are irreducible for the Hecke algebra $H_n(q)$ when q is specialized to a primitive ℓ^{th} root of unity. An ℓ -regular partition λ indexes a simple module D^λ for $H_n(q)$ when q is a primitive ℓ^{th} root of unity. A weak ℓ -partition λ is an ℓ -regular partition for which there is a (not necessarily ℓ -regular) partition μ so that $S^\mu = D^\lambda$. We noticed that within the crystal $B(\Lambda_0)$ the weak ℓ -partitions satisfied rules similar to the rules given in [1] for ℓ -partitions. In order to prove

Supported in part by NSF grant DMS-0135345.

this, we built a modified version of the crystal $B(\Lambda_0)$, which we denote $B(\Lambda_0)^L$. The combinatorial description of $B(\Lambda_0)^L$ satisfies the following properties:

- The nodes of $B(\Lambda_0)^L$ are partitions, and there is an i -arrow from λ to μ only when the difference $\mu \setminus \lambda$ is a box of residue i .
- If λ is a node of $B(\Lambda_0)$ and is a weak ℓ -partition with $D^\lambda = S^\nu$ for some partition ν then ν is a node of $B(\Lambda_0)^L$.
- $B(\Lambda_0) \cong B(\Lambda_0)^L$ and this crystal isomorphism yields an interesting bijection on the nodes. The map being used for the isomorphism has been well studied [10], but never before in the context of a crystal isomorphism.
- There exists elementary combinatorial arguments which generalize the crystal theoretic results of $B(\Lambda_0)$ to $B(\Lambda_0)^L$.
- The partitions which are nodes of $B(\Lambda_0)^L$ can be identified by a simple combinatorial condition.

The new description of the crystal $B(\Lambda_0)$ is in many ways more important than the theorems which were proven by the existence of it. Besides the fact that it is a useful tool in proving theorems about $B(\Lambda_0)$, our new description also gives a new set of partitions (in bijection to ℓ -regular partitions), which can be interpreted in terms of the representation theory of $H_n(q)$.

1.1. Background and Previous Results. Let λ be a partition of n (written $\lambda \vdash n$) and $\ell \geq 3$ be an integer. We will use the convention (x, y) to denote the box which sits in the x^{th} row and the y^{th} column of the Young diagram of λ . We denote the transpose of λ by λ' . Throughout this paper, all of our partitions are drawn in English notation. Sometimes the shorthand (a^k) will be used to represent the rectangular partition which has k -parts, all of size a . $\mathcal{P}$ will denote the set of all partitions. An ℓ -regular partition is one in which no part occurs ℓ or more times. The *length of a partition* λ will be the number of nonzero parts of λ and will be denoted $\text{len}(\lambda)$.

The *hook length* of the (a, c) box of λ is defined to be the number of boxes to the right and below the box (a, c) , including the box (a, c) itself. It will be denoted $h_{(a, c)}^\lambda$. The *arm* of the (a, c) box of λ is defined to be the number of boxes to the right of the box (a, c) , *not* including the box (a, c) . It will be denoted $\text{arm}(a, c)$.


Similarly, the leg is all boxes below (a, c) , not including (a, c) and will be denoted $leg(a, c)$.

A removable ℓ -rim hook in λ is a connected sequence of ℓ boxes in the Young diagram of λ , containing no 2×2 square, such that when removed from λ , the remaining diagram is the Young diagram of some other partition.


Any partition which has no removable rim hooks of length ℓ is called an ℓ -core. Equivalently, λ is an ℓ -core if for every box $(i, j) \in \lambda$, $\ell \nmid h_{(i, j)}^\lambda$. The set of all ℓ -cores will be denoted $\mathcal{C}_\ell$. Any partition λ has an ℓ -core, which is obtained by removing ℓ -rim hooks from the outer edge while at each step the removal of a hook is still a (non-skew) partition. A core is uniquely determined from the partition, independently of choice of successively removing rim hooks. The number of ℓ -rim hooks which must be removed from a partition λ to obtain its core is called the ℓ -weight of λ (or simply the weight of λ when ℓ is clear from the context). See [10] for more details.

Removable ℓ -rim hooks which are flat (whose boxes are contained in one row of a partition) will be called *horizontal ℓ -rim hooks*. Removable ℓ -rim hooks which are not flat will be called *non-horizontal ℓ -rim hooks*. A removable ℓ -rim hook contained entirely in a single column of the Young diagram of a partition will be called a *vertical ℓ -rim hook*. Removable ℓ -rim hooks not contained in a single column will be called *non-vertical ℓ -rim hooks*. Two connected sequences of boxes will be called *adjacent* if there exists boxes in each which share an edge.

Example 1.1.1. Let $\lambda = (3, 2, 1)$ and let $\ell = 3$. Then the boxes $(1, 2)$, $(1, 3)$ and $(2, 2)$ comprise a (non-vertical, non-horizontal) 3-rim hook. After removal of this 3-rim hook, the remaining partition is $(1, 1, 1)$, which is a vertical 3-rim hook. Hence the 3-core of λ is the empty partition. These two 3-rim hooks are adjacent.




Example 1.1.2. Let $\lambda = (4, 1, 1, 1)$ and $\ell = 3$. Then λ has two 3-rim hooks (one horizontal and one vertical). They are not adjacent.




Definition 1.1.3. An ℓ -partition is an ℓ -regular partition with no removable non-horizontal ℓ -rim hooks, such that after removing any number of horizontal ℓ -rim hooks, the remaining diagram still has no removable non-horizontal ℓ -rim hooks.

Example 1.1.4. Any ℓ -core is an ℓ -partition.

Example 1.1.5. $(6, 4)$ is a 3-partition, as well as a ℓ -partition for $\ell > 7$ but is not a 2, 4, 5, 6, 7-partition



Example 1.1.6. $(5, 4, 1)$ is a 6-core. It is a 2-partition, but not a 2-core. It is not a 3, 4, 5, 7-partition. It is an ℓ -core for $\ell > 7$.



We introduce the following algebra to explain some of the representation theoretic significance of the combinatorics described in this paper.

Definition 1.1.7. For a fixed field $\mathbb{F}$ of characteristic zero and $0 \neq q \in \mathbb{F}$, the finite Hecke Algebra $H_n(q)$ is defined to be the $\mathbb{F}$ -algebra generated by $T_1, \dots, T_{n-1}$ with relations

$$\begin{aligned} T_i T_j &= T_j T_i && \text{for } |i - j| > 1 \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1} && \text{for } 1 \leq i < n - 1 \\ T_i^2 &= (q - 1)T_i + q && \text{for } 1 \leq i \leq n - 1. \end{aligned}$$

In this paper we will always assume that $q \neq 1$ and that $q \in \mathbb{F}$ is a primitive ℓ^{th} root of unity in a field $\mathbb{F}$ of characteristic zero.

Remark 1.1.8. When q is specialized to 1 the Hecke algebra becomes the group algebra of the symmetric group.

Similar to the symmetric group, a construction of the Specht module $S^\lambda = S^\lambda[q]$ exists for $H_n(q)$ (see [3]). Let ℓ be an integer greater than 1. Let

$$\nu_\ell(k) = \begin{cases} 1 & \ell \mid k \\ 0 & \ell \nmid k. \end{cases}$$

It is known that over the finite Hecke algebra $H_n(q)$, when q is a primitive ℓ^{th} root of unity, the Specht module S^λ for an ℓ -regular partition λ is irreducible if and only if

$$(\star) \quad \nu_\ell(h_{(a,c)}^\lambda) = \nu_\ell(h_{(b,c)}^\lambda) \quad \text{for all pairs } (a, c), (b, c) \in \lambda$$

(see [11]). So, an equivalent condition for the irreducibility of the Specht module indexed by an ℓ -regular partition is that the hook lengths in every column of the partition λ are either all divisible by ℓ or none of them are. In [1], we proved the following.

Theorem 1.1.9. *A partition is an ℓ -partition if and only if it is ℓ -regular and satisfies $(\star)$.*

Fayers recently settled the following conjecture of James and Mathas.

Theorem 1.1.10 (Fayers, [4]). *Suppose $\ell > 2$. Let λ be a partition. Then S^λ is reducible if and only if there exists boxes (a, b) (a, y) and (x, b) in the Young diagram of λ for which:*

- $\nu_\ell(h_{(a,b)}^\lambda) = 1$
- $\nu_\ell(h_{(a,y)}^\lambda) = \nu_\ell(h_{(x,b)}^\lambda) = 0$

A partition which has no such boxes is called an $(\ell, 0)$ -JM partition. Equivalently, λ is an $(\ell, 0)$ -JM partition if and only if the Specht module S^λ is irreducible.

All of the irreducible representations of $H_n(q)$ have been constructed when q is a primitive ℓ^{th} root of unity. These modules are indexed by ℓ -regular partitions λ , and are called D^λ . D^λ is the unique simple quotient of S^λ (see [3] for more details). In particular $D^\lambda = S^\lambda$ if and only if S^λ is irreducible and λ is ℓ -regular. For λ not necessarily ℓ -regular, S^λ is irreducible if and only if there exists an ℓ -regular partition μ so that $S^\lambda \cong D^\mu$.

1.2. Summary of results from this paper. In Section 2 we give a new way of characterizing $(\ell, 0)$ -JM partitions by their removable ℓ -rim hooks. In Section 3 we use this result to count the number of $(\ell, 0)$ -JM partitions in a block. In Section 4 we recall the description of the crystal $B(\Lambda_0)$ of $\widehat{\mathfrak{sl}}_\ell$ involving ℓ -regular partitions. In Section 5 we give our new description of the crystal $B(\Lambda_0)$. Section 6 is a brief introduction to the map on partitions known as regularization, which was first described by James. Section 7 gives a new procedure for finding an inverse for the map of regularization. Section 8 extends our crystal theorems from [1] to the crystal $B(\Lambda_0)^L$. Section 9 reinterprets the classical crystal rules in the combinatorial framework of the new crystal rules. Section 10 contains the proof that the two descriptions of the crystal $B(\Lambda_0)$ are the same; the isomorphism being regularization. Section 11 transfers the crystal theorems on the new crystal to theorems on the old crystal by using this isomorphism. In Section 12 we give a purely representation theoretic proof of one of our main theorems (Theorem 4.3.5). Lastly, in Section 13 we show some connections between the ladder crystal and the Mullineux map, which leads us to a nice classification of the partitions in $B(\Lambda_0)^L$.

1.3. Acknowledgments. The author would like to thank several people who made this paper possible. I first showed the idea of the ladder crystal to Steve Pon in the summer of 2007 and with him we sketched out some work that would later be the basis for proving that the two crystals were isomorphic. Later on, I ran into a problem showing the existence of a smallest partition in dominance order in each regularization class and together with Brant Jones we developed an algorithm for finding such a partition, which I've generalized in this paper in Section 7. After sending a draft to Matthew Fayers, he responded with helpful suggestions and corrections. His comments also lead me to write Section 13. Finally, I would like to thank my advisor Monica Vazirani. Finding answers to her questions led me to discover the ladder crystal, and numerous ensuing discussions helped in the coherency and direction of this paper.

2. CLASSIFYING $(\ell, 0)$ -JM PARTITIONS BY THEIR REMOVABLE ℓ -RIM HOOKS

2.1. Motivation. In this section we give a new description of $(\ell, 0)$ -JM partitions. This condition is related to how ℓ -rim hooks are removed from a partition and

is a generalization of Theorem 2.1.6 in [1] about ℓ -partitions. The condition we give (equivalent to the condition conjectured by James and Mathas and proved by Fayers) will be used in several proofs throughout this paper.

2.2. Removing ℓ -Rim Hooks and $(\ell, 0)$ -JM partitions.

Definition 2.2.1. Let λ be a partition. Let $\ell > 2$. Then λ is a *generalized ℓ -partition* if:

- (1) It has no removable ℓ -rim hooks other than horizontal and vertical ℓ -rim hooks.
- (2) If R is a vertical (resp. horizontal) ℓ -rim hook of λ and S is a horizontal (resp. vertical) ℓ -rim hook of $\lambda \setminus R$, then R and S are not adjacent.
- (3) After removing any sequence of horizontal and vertical ℓ -rim hooks from the Young diagram of λ , the remaining partition satisfies (1) and (2).

Example 2.2.2. Let $\lambda = (3, 1, 1, 1)$. Then λ is not a generalized 3-partition. λ has a vertical 3-rim hook R containing the boxes $(2, 1), (3, 1), (4, 1)$. Removing R leaves a horizontal 3-rim hook S containing the boxes $(1, 1), (1, 2), (1, 3)$. S is adjacent to R , so λ is not a generalized 3-partition.

S	S	S
R		
R		
R		

Remark 2.2.3. We will sometimes abuse notation and say that R and S in Example 2.2.2 are adjacent vertical and horizontal ℓ -rim hooks. The meaning here is not that they are both ℓ -rim hooks of λ (S is not an ℓ -rim hook of λ), but rather that they are an example of a violation of condition 2 from Definition 2.2.1

We will need a few lemmas before we come to our main theorem of this section, which states that the notions of $(\ell, 0)$ -JM partitions and generalized ℓ -partitions are equivalent.

Lemma 2.2.4. *Suppose λ is not an $(\ell, 0)$ -JM partition. Then a partition obtained from λ by adding a horizontal or vertical ℓ -rim hook is also not an $(\ell, 0)$ -JM partition.*

The proof is nearly identical to our lemma in [1] about adding horizontal ℓ -rim hooks to an ℓ -regular $(\ell, 0)$ -JM partition, so we refer the reader there.

Then next lemma gives an equivalent definition of an ℓ -core described in terms of hook lengths.

Lemma 2.2.5. *A partition λ is an ℓ -core if and only if the hook length $h_{(a,b)}^\lambda$ is not divisible by ℓ for every box (a,b) in the Young diagram of λ .*

The upcoming lemma simplifies the condition for being an $(\ell, 0)$ -JM partition and is used in the proof of Theorem 2.2.7.

Lemma 2.2.6. *Suppose λ is not an $(\ell, 0)$ -JM partition. Then there exists boxes (c, d) , (c, w) and (z, d) with $c < z$, $d < w$, and $\ell \mid h_{(c,d)}^\lambda$, $\ell \nmid h_{(c,w)}^\lambda, h_{(z,d)}^\lambda$.*

Proof. The condition proven by Fayers [4] says that there exists boxes (a, b) , (a, y) and (x, b) where $\ell \mid h_{(a,b)}^\lambda$ and $\ell \nmid h_{(a,y)}^\lambda, h_{(x,b)}^\lambda$. If $a < x$ and $b < y$ then we are done. The other cases follow below:

Case 1: That $x < a$ and $y < b$. Assume no triple exists satisfying the statement of the lemma. Then either all boxes to the right of the (a, b) box will have hook lengths divisible by ℓ , or all boxes below will. Without loss of generality, suppose that all boxes below the (a, b) box have hook lengths divisible by ℓ . Let $c < a$ be the largest integer so that $\ell \nmid h_{(c,b)}$. Let $z = c + 1$. Then one of the boxes $(c, b+1), (c, b+2), \dots, (c, b+\ell-1)$ has a hook length divisible by ℓ . This is because the box (h, b) at the bottom of column b has a hook length divisible by ℓ , so the hook lengths $h_{(c,b)}^\lambda = h_{(c,b+1)}^\lambda + 1 = \dots = h_{(c,b+\ell-1)}^\lambda + \ell - 1$. Suppose it is (c, d) . Then $\ell \nmid h_{(z,d)}^\lambda$ since $h_{(z,b)} = h_{(z,d)} + d - b$ and $d - b < \ell$.

If $d \neq b + \ell - 1$ or $h_{(h,b)}^\lambda > \ell$ then letting $w = d + 1$ gives (c, w) to the right of (c, d) so that $\ell \nmid h_{(c,w)}^\lambda$ (in fact $h_{(c,w)}^\lambda = h_{(c,d)}^\lambda - 1$).

If $d = b + \ell - 1$ and $h_{(h,b)}^\lambda = \ell$ then there is a box in spot $(c, d + 1)$ with hook length $h_{(c,d+1)}^\lambda = h_{(c,d)}^\lambda - 2$ since there must be a box in the position $(h - 1, d + 1)$, due to the fact that $\ell \mid h_{(h-1,b)}^\lambda$ and $h_{(h-1,b)}^\lambda > \ell$. Letting $w = d + 1$ again yields $\ell \nmid h_{(c,w)}^\lambda$. Note that this requires that $\ell > 2$. In fact if $\ell = 2$ we cannot even be sure that there is a box in position $(c, d + 1)$.

Case 2: That $x < a$ and $y > b$. If there was a box (n, b) ($n > a$) with a hook length not divisible by ℓ then we would be done. So we can assume that all hook

lengths in column b below row a are divisible by ℓ . Let $c < a$ be the largest integer so that $\ell \nmid h_{(c,b)}^\lambda$. Let $z = c+1$. Similar to Case 1 above, we find a d so that $\ell \mid h_{(c,d)}^\lambda$. Then $\ell \nmid h_{(z,d)}^\lambda$ and by the same argument as in Case 1, if we let $w = d+1$ then $\ell \nmid h_{(c,w)}^\lambda$.

Case 3: That $x > a$ and $y < b$. Then transpose λ , apply Case 2, and transpose back. $\square$

Theorem 2.2.7. *A partition is an $(\ell, 0)$ -JM partition if and only if it is a generalized ℓ -partition.*

Proof. Suppose that λ is not a generalized ℓ -partition. Then remove non-adjacent horizontal and vertical ℓ -rim hooks until you obtain a partition μ which has either a non-vertical non-horizontal ℓ -rim hook, or adjacent horizontal and vertical ℓ -rim hooks. If there is a non-horizontal, non-vertical ℓ -rim hook in μ , let's say the ℓ -rim hook has southwest most box (a, b) and northeast most box (c, d) . Then $\ell \mid h_{(c,b)}^\mu$ but $\ell \nmid h_{(a,b)}^\mu, h_{(c,d)}^\mu$ since $h_{(a,b)}^\mu = h_{(c,d)}^\mu = 1$. We know $a \neq c$ and $b \neq d$ since our ℓ -rim hook is non-horizontal and non-vertical. Therefore, μ is not an $(\ell, 0)$ -JM partition. By Lemma 2.2.4, λ is not an $(\ell, 0)$ -JM partition. Similarly, if μ has adjacent vertical and horizontal ℓ -rim hooks, then let (a, b) be the southwest most box in the vertical ℓ -rim hook and let (c, d) be the position of the northeast most box in the horizontal ℓ -rim hook. Again, $\ell \mid h_{(c,b)}^\mu$ but $\ell \nmid h_{(a,b)}^\mu, h_{(c,d)}^\mu$. Therefore μ cannot be an $(\ell, 0)$ -JM partition, so λ is not an $(\ell, 0)$ -JM partition.

Conversely, suppose λ is not an $(\ell, 0)$ -JM partition, but that it is a generalized ℓ -partition. Then by Lemma 2.2.6 there are boxes (a, b) , (a, y) and (x, b) with $a < x$ and $b < y$, which satisfy $\ell \mid h_{(a,b)}^\lambda$, and $\ell \nmid h_{(a,y)}^\lambda, h_{(x,b)}^\lambda$. Form a new partition μ by taking all of the boxes (m, n) in λ such that $m \geq a$ and $n \geq b$. Since λ was a generalized ℓ -partition, μ must be also. By Lemma 2.2.5, we know that there must exist a removable ℓ -rim hook from μ , since $\ell \mid h_{(1,1)}^\mu$. Since μ is a generalized ℓ -partition, the ℓ -rim hook must be either horizontal or vertical. If we remove a horizontal ℓ -rim hook we can only change $h_{(1,1)}^\mu$ by either 0, 1 or ℓ .

If it was changed by 1, then we let $(x', 1)$ denote the box in μ which corresponds to (x, b) in λ . Letting ν denote the partition μ with the horizontal ℓ -rim hook removed, we then note that $\ell \mid h_{(1,\ell)}^\nu$, $\ell \nmid h_{(1,1)}^\nu u$ and $\ell \nmid h_{(x',\ell)}^\nu u$. This is because $h_{(1,\ell)}^\mu = 1 \pmod{\ell}$, $h_{(1,1)}^\mu u = 0 \pmod{\ell}$, $h_{(x',\ell)}^\mu = h_{(x',1)}^\mu + 1 \pmod{\ell}$ and the fact that

removing the ℓ -rim hook question will drop each of these hook lengths by exactly 1. Hence we deduce that ν is not an $(\ell, 0)$ -JM partition, but it is still a generalized ℓ -partition. In this case we start this process over, with ν in place of λ . We do the same for vertical ℓ -rim hooks which change the hook length $h_{(1,1)}^\mu$ by 1.

Now we may assume that removing horizontal or vertical ℓ -rim hooks from μ will not change that ℓ divides $h_{(1,1)}^\mu$ (because removing each ℓ -rim hook will change the hook length $h_{(1,1)}^\mu$ by either 0 or ℓ). Therefore we can keep removing ℓ -rim hooks until we have removed box $(1,1)$ entirely, in which case the remaining partition had a horizontal ℓ -rim hook adjacent to a vertical ℓ -rim hook (since both (x, b) and (a, y) must have been removed, the ℓ -rim hooks could not have been exclusively horizontal or vertical). This contradicts μ being a generalized ℓ -partition. $\square$

Example 2.2.8. Let $\lambda = (10, 8, 3, 2^2, 1^5)$. Then λ is a generalized 3-partition and a $(3, 0)$ -JM partition. λ is drawn below with each hook length $h_{(a,b)}^\lambda$ written in the box (a, b) and the possible removable ℓ -rim hooks outlined. Also, hook lengths which are divisible by ℓ are underlined.

19	13	10	8	7	<u>6</u>	5	4	2	1
16	10	7	5	4	<u>3</u>	2	1		
10	4	1							
8	2								
7	1								
5									
4									
<u>3</u>									
2									
1									

Example 2.2.9. $\lambda = (3, 1, 1, 1)$ from Example 2.2.2 is not a $(3, 0)$ -JM partition and it is not a generalized 3-partition.

3. DECOMPOSITION OF $(\ell, 0)$ -JM PARTITIONS

3.1. Motivation. In [1] we gave a decomposition of ℓ -partitions. In this section we give a similar decomposition for all $(\ell, 0)$ -JM partitions. This decomposition is

important for the proofs of the theorems in later sections. It also implies Theorem 3.2.3, which counts all $(\ell, 0)$ -JM partitions of a given core and weight.

3.2. Decomposing $(\ell, 0)$ -JM partitions. Let μ be an ℓ -core with $\mu_1 - \mu_2 < \ell - 1$ and $\mu'_1 - \mu'_2 < \ell - 1$. Let $r, s \geq 0$. Let ρ and σ be partitions with $\text{len}(\rho) \leq r + 1$ and $\text{len}(\sigma) \leq s + 1$. If $\mu = \emptyset$ then we require at least one of ρ_{r+1}, σ_{s+1} to be zero. Following the construction of [1], we construct a partition corresponding to $(\mu, r, s, \rho, \sigma)$ as follows. Starting with μ , attach r rows above μ , with each row $\ell - 1$ boxes longer than the previous. Then attach s columns to the left of μ , with each column $\ell - 1$ boxes longer than the previous. This partition will be denoted $(\mu, r, s, \emptyset, \emptyset)$. Formally, if $\mu = (\mu_1, \mu_2, \dots, \mu_m)$ then $(\mu, r, s, \emptyset, \emptyset)$ represents the partition (which is an ℓ -core):

$$(s + \mu_1 + r(\ell - 1), s + \mu_1 + (r - 1)(\ell - 1), \dots, s + \mu_1 + \ell - 1, s + \mu_1,$$

$$s + \mu_2, \dots, s + \mu_m, s^{\ell-1}, (s - 1)^{\ell-1}, \dots, 1^{\ell-1})$$

where $s^{\ell-1}$ stands for $\ell - 1$ copies of s . Now to the first $r + 1$ rows attach ρ_i horizontal ℓ -rim hooks to row i . Similarly, to the first $s + 1$ columns, attach σ_j vertical ℓ -rim hooks to column j . The resulting partition λ corresponding to $(\mu, r, s, \rho, \sigma)$ will be


$$\lambda = (s + \mu_1 + r(\ell - 1) + \rho_1\ell, s + \mu_1 + (r - 1)(\ell - 1) + \rho_2\ell, \dots,$$

$$s + \mu_1 + (\ell - 1) + \rho_r\ell, s + \mu_1 + \rho_{r+1}\ell, s + \mu_2, s + \mu_3, \dots,$$


$$s + \mu_m, (s + 1)^{\sigma_{s+1}\ell}, s^{\ell-1+(\sigma_s-\sigma_{s+1})\ell}, (s - 1)^{\ell-1+(\sigma_{s-1}-\sigma_s)\ell}, \dots, 1^{\ell-1+(\sigma_1-\sigma_2)\ell}).$$

We denote this decomposition as $\lambda \approx (\mu, r, s, \rho, \sigma)$.

Example 3.2.1. Let $\ell = 3$ and $(\mu, r, s, \rho, \sigma) = ((1), 3, 2, (2, 1, 1, 1), (2, 1, 0))$. Then $((1), 3, 2, \emptyset, \emptyset)$ is drawn below, with μ framed.



$((1), 3, 2, (2, 1, 1, 1), (2, 1, 0))$ is drawn below, now with $((1), 3, 2, \emptyset, \emptyset)$ framed.



Theorem 3.2.2. *If $\lambda \approx (\mu, r, s, \rho, \sigma)$ (with at least one of $\rho_{r+1}, \sigma_{s+1} = 0$ if $\mu = \emptyset$), then λ is an $(\ell, 0)$ -JM partition. Conversely, all such $(\ell, 0)$ -JM partitions are of this form.*

Proof. If $\lambda \approx (\mu, r, s, \rho, \sigma)$ then it is clear by construction that λ satisfies the criterion for a generalized ℓ -partition (see Definition 2.2.1). By Theorem 2.2.7, λ is an $(\ell, 0)$ -JM partition.

Conversely, if λ is an $(\ell, 0)$ -JM partition then by Theorem 2.2.7 its only removable ℓ -rim hooks are horizontal or vertical. Let ρ_i be the number of removable horizontal

ℓ -rim hooks in row i which are removed in going to the ℓ -core of λ , and let σ_j be the number of removable vertical ℓ -rim hooks in column j (since λ has no adjacent ℓ -rim hooks, these numbers are well defined). Once all ℓ -rim hooks are removed, let r (resp. s) be the number of rows (resp. columns) whose successive differences are $\ell - 1$. Removing these topmost r rows and leftmost s columns leaves an ℓ -core μ . Then $\lambda \approx (\mu, r, s, \rho, \sigma)$. $\square$

Further in the text, we will make use of Theorem 3.2.2. Many times we will show that a partition λ is an $(\ell, 0)$ -JM partition by giving an explicit decomposition of λ into $(\mu, r, s, \rho, \sigma)$.

The following theorem follows directly from Theorem 3.2.2. We have decided to include it after reading [2] and recognizing the similarity between their construction of the (ℓ, ℓ) -JM partitions and our construction. Let $\Gamma_\ell(\nu) = \{\lambda : \lambda \text{ is an } (\ell, 0)\text{-JM partition with core } \nu\}$.

Theorem 3.2.3. *Suppose ν is an ℓ -core, with*

- $\nu_1 - \nu_2 = \nu_2 - \nu_3 = \cdots = \nu_r - \nu_{r+1} = \ell - 1$ and $\nu_{r+1} - \nu_{r+2} \neq \ell - 1$,
- $\nu'_1 - \nu'_2 = \nu'_2 - \nu'_3 = \cdots = \nu'_s - \nu'_{s+1} = \ell - 1$ and $\nu'_{s+1} - \nu'_{s+2} \neq \ell - 1$.

If ν also satisfies $\nu_{r+1} > s$ then the number of $(\ell, 0)$ -JM partitions of a fixed ℓ -weight w with core ν , denoted $\Gamma_\ell(\nu)$, is the number of pairs of partitions (α, β) , $\text{len}(\alpha) \leq r+1$, $\text{len}(\beta) \leq s+1$ whose size sums to w ($|\alpha| + |\beta| = w$). The generating function for all $(\ell, 0)$ -JM partitions with core ν with respect to the statistic of the ℓ -weight of the partition is thus $\prod_{i=0}^{r+1} \frac{1}{1-x^i} \prod_{j=0}^{s+1} \frac{1}{1-x^j}$. Hence the generating function for all $(\ell, 0)$ -JM partitions for a fixed core ν as above with respect to the statistic of the size of the partition is

$$\sum_{\lambda \in \Gamma_\ell(\nu)} x^{|\lambda|} = x^{|\nu|} \prod_{i=0}^{r+1} \frac{1}{1-x^{\ell i}} \prod_{j=0}^{s+1} \frac{1}{1-x^{\ell j}}.$$

If ν satisfies $\nu_{r+1} = s$ then the number of $(\ell, 0)$ -JM partitions of a fixed ℓ -weight w with core ν is the number of pairs of partitions (α, β) , $\text{len}(\alpha) \leq r+1$, $\text{len}(\beta) \leq s$ whose sizes sum to w plus the number of pairs of partitions (γ, δ) , $\text{len}(\gamma) \leq r$, $\text{len}(\delta) \leq s+1$ whose sizes sum to w . The generating function for all $(\ell, 0)$ -JM partitions with core ν with respect to the statistic of the ℓ -weight of the partition is

thus $\left(\frac{1}{1-x^{r+1}} + \frac{1}{1-x^{s+1}}\right) \prod_{i=0}^r \frac{1}{1-x^i} \prod_{j=0}^s \frac{1}{1-x^j}$. Hence the generating function for all $(\ell, 0)$ -JM partitions for a fixed core ν with respect to the statistic of the size of the partition is

$$\sum_{\lambda \in \Gamma_\ell(\nu)} x^{|\lambda|} = x^{|\nu|} \left(\frac{1}{1-x^{\ell(r+1)}} + \frac{1}{1-x^{\ell(s+1)}} \right) \prod_{i=0}^r \frac{1}{1-x^{\ell i}} \prod_{j=0}^s \frac{1}{1-x^{\ell j}}.$$

Example 3.2.4. Let $\ell = 3$ and $\lambda = (3, 1)$. Then λ is a 3-core. λ has $r = 1$ and $s = 0$. Suppose we are looking for the number of $(3, 0)$ -JM partitions with core λ and size 13 (i.e. weight 3). There are as many such $(3, 0)$ -JM partitions as there are pairs of partitions (α, β) with $\text{len}(\alpha) \leq 2$, $\text{len}(\beta) \leq 1$ and $|\alpha| + |\beta| = 3$. There are 6 such pairs of partitions. They are $((3), \emptyset), ((2, 1), \emptyset), ((2), (1)), ((1, 1), (1)), ((1), (2))$ and $(\emptyset, (3))$. They correspond to the 6 $(3, 0)$ -JM partitions $(12, 1)$, $(9, 4)$, $(9, 1^4)$, $(6, 4, 1^3)$, $(6, 1^7)$ and $(3, 1^{10})$ respectively.

4. CLASSICAL DESCRIPTION OF $B(\Lambda_0)$

4.1. Introduction. In this section, we recall the description of the crystal graph $B(\Lambda_0)$ currently used in the literature, first described by Misra and Miwa [18]. We end this section with the results from [1] which we will generalize.

4.2. Classical description of the crystal $B(\Lambda_0)$. We will assume some familiarity with the theory of crystals (see [12] for details), and their relationship to the representation theory of the finite Hecke Algebra (see [7], [13] for details). We will look at the crystal $B(\Lambda_0)$ of the irreducible highest weight module $V(\Lambda_0)$ of the affine Lie algebra $\widehat{\mathfrak{sl}_\ell}$ (also called the basic representation of $\widehat{\mathfrak{sl}_\ell}$). The nodes of $B(\Lambda_0)$ are ℓ -regular partitions. The set of nodes will be denoted $B := \{\lambda \in \mathcal{P} : \lambda \text{ is } \ell\text{-regular}\}$. We will describe the arrows of $B(\Lambda_0)$ below.

We view the Young diagram for λ as a set of boxes, with their corresponding residues $b - a \pmod{\ell}$ written into the box (a, b) . A box in λ is said to be a removable i -box if it has residue i and after removing that box the remaining diagram is still a partition. A space not in λ is an addable i -box if it has residue i and adding that box to λ yields a partition.

Example 4.2.1. Let $\lambda = (8, 5, 4, 1)$ and $\ell = 3$. Then the residues are filled into the corresponding Young diagram as follows:

$$\lambda = \begin{array}{cccccccc} 0 & 1 & 2 & 0 & 1 & 2 & 0 & 1 \\ 2 & 0 & 1 & 2 & 0 & 1 & & \\ 1 & 2 & 0 & 1 & 2 & & & \\ 0 & 1 & & & & & & \end{array}$$

Here λ has two removable 0-boxes (boxes (2,5) and (4,1)), two removable 1-boxes (boxes (1,8) and (3,4)), no removable 2-boxes, no addable 0-boxes, two addable 1-boxes (in positions (2,6) and (4,2)), and three addable 2-boxes (in positions (1,9), (3,5) and (5,1)).

For a fixed i , ($0 \leq i < \ell$), we place $-$ in each removable i -box and $+$ in each addable i -box. The i -signature of λ is the word of $+$ and $-$'s in the diagram for λ , written from bottom left to top right. The reduced i -signature is the word obtained after repeatedly removing from the i -signature all adjacent pairs $-+$. The resulting word will now be of the form $+\cdots+++-\cdots-$. The boxes corresponding to $-$'s in the reduced i -signature are called *normal i -boxes*, and the boxes corresponding to $+$'s are called *conormal i -boxes*. $\varepsilon_i(\lambda)$ is defined to be the number of normal i -boxes of λ , and $\varphi_i(\lambda)$ is defined to be the number of conormal i -boxes. If a leftmost $-$ exists, the box corresponding to such a $-$ is called the *good i -box* of λ . If a rightmost $+$ exists, the box corresponding to such a $+$ is called the *cogood i -box*. All of these definitions can be found in Kleshchev's book [13].

Example 4.2.2. Let $\lambda = (8, 5, 4, 1)$ and $\ell = 3$ be as above. Fix $i = 1$. The diagram for λ with removable and addable 1-boxes marked looks like:

								-
						+		
				-				
	+							

The 1-signature of λ is $+-+ -$, so the reduced 1-signature is $+-$ and the diagram has a good 1-box in the first row, and a cogood 1-box in the fourth row. Here $\varepsilon_1(\lambda) = 1$ and $\varphi_1(\lambda) = 1$.

We recall the action of the crystal operators on B . The crystal operator $\tilde{e}_i : B \xrightarrow{i} B \cup \{0\}$ assigns to a partition λ the partition $\tilde{e}_i(\lambda) = \lambda \setminus x$, where x is




FIGURE 1. The 1-string of $(7, 5, 4, 1)$ in $B(\Lambda_0)$ for $\ell = 3$ from Example 4.2.3.

the good i -box of λ . If no such box exists, then $\tilde{e}_i(\lambda) = 0$. It is clear then that $\varepsilon_i(\lambda) = \max\{k : \tilde{e}_i^k \lambda \neq 0\}$.

Similarly, $\tilde{f}_i : B \xrightarrow{i} B \cup \{0\}$ is the operator which assigns to a partition λ the partition $\tilde{f}_i(\lambda) = \lambda \cup x$, where x is the cogood i -box of λ . If no such box exists, then $\tilde{f}_i(\lambda) = 0$. It is clear then that $\varphi_i(\lambda) = \max\{k : \tilde{f}_i^k \lambda \neq 0\}$.

For $i \in \mathbb{Z}/\ell\mathbb{Z}$, we write $\lambda \xrightarrow{i} \mu$ to stand for $\tilde{f}_i \lambda = \mu$. We say that there is an i -arrow from λ to μ . Note that $\lambda \xrightarrow{i} \mu$ if and only if $\tilde{e}_i \mu = \lambda$. A maximal chain of consecutive i -arrows will be called an i -string. We note that the empty partition $\emptyset$ is the unique highest weight node of the crystal (i.e. $\tilde{e}_i \emptyset = 0$ for every $i \in \mathbb{Z}/\ell\mathbb{Z}$.) For a picture of a part of this crystal graph, see [14] for the cases $\ell = 2$ and 3.

Example 4.2.3. Continuing with the above example, we see that $\tilde{e}_1(8, 5, 4, 1) = (7, 5, 4, 1)$ and $\tilde{f}_1(8, 5, 4, 1) = (8, 5, 4, 2)$. Also, $\tilde{e}_1^2(8, 5, 4, 1) = 0$ and $\tilde{f}_1^2(8, 5, 4, 1) = 0$. The sequence $(7, 5, 4, 1) \xrightarrow{1} (8, 5, 4, 1) \xrightarrow{1} (8, 5, 4, 2)$ is a 1-string of length 3.

For the rest of this paper, $\varphi = \varphi_i(\lambda)$ and $\varepsilon = \varepsilon_i(\lambda)$.

4.3. Previous results on the crystal $B(\Lambda_0)$. In [1] we proved the following theorems about ℓ -partitions in the crystal $B(\Lambda_0)$.

Theorem 4.3.1. *Suppose that λ is an ℓ -partition and $0 \leq i < \ell$. Then*

- (1) $\tilde{f}_i^\varphi \lambda$ is an ℓ -partition,
- (2) $\tilde{e}_i^\varepsilon \lambda$ is an ℓ -partition.

Theorem 4.3.2. *Suppose that λ is an ℓ -partition. Then*

- (1) $\tilde{f}_i^k \lambda$ is not an ℓ -partition for $0 < k < \varphi - 1$,
- (2) $\tilde{e}_i^k \lambda$ is not an ℓ -partition for $1 < k < \varepsilon$.

Definition 4.3.3. An ℓ -regular partition λ will be called a *weak ℓ -partition* if there exists a partition μ so that $S^\mu = D^\lambda$.

Remark 4.3.4. We note that all ℓ -partitions are trivially weak ℓ -partitions (since $S^\lambda = D^\lambda$ if λ is an ℓ -partition). Also, all weak ℓ -partitions are by definition ℓ -regular.

In this paper, we generalize the above Theorems 4.3.1 and 4.3.2 to weak ℓ -partitions. We first give the statements of our new theorems.

Theorem 4.3.5. *Suppose that λ is a weak ℓ -partition and $0 \leq i < \ell$. Then*

- (1) $\tilde{f}_i^\varphi \lambda$ is a weak ℓ -partition,
- (2) $\tilde{e}_i^\varepsilon \lambda$ is a weak ℓ -partition.

Theorem 4.3.6. *Suppose that λ is a weak ℓ -partition. Then*

- (1) $\tilde{f}_i^k \lambda$ is not a weak ℓ -partition for $0 < k < \varphi - 1$,
- (2) $\tilde{e}_i^k \lambda$ is not a weak ℓ -partition for $1 < k < \varepsilon$.


To prove the above theorems, we start by introducing our new description of the crystal $B(\Lambda_0)$.

5. THE LADDER CRYSTAL: $B(\Lambda_0)^L$

5.1. Ladders. We first recall what a ladder is in regards to a partition. Let λ be a partition and let $\ell > 2$ be a fixed integer. For any box (a, b) in the Young diagram of λ , the *ladder* of (a, b) is the set of all positions (c, d) which satisfy $\frac{c-a}{d-b} = \ell - 1$ and $c, d > 0$.

Remark 5.1.1. The definition implies that two boxes in the same ladder will share the same residue. An i -ladder will be a ladder which has residue i .

Example 5.1.2. Let $\lambda = (3, 3, 1)$, $\ell = 3$. Then there is a 1-ladder which contains the boxes $(1, 2)$ and $(3, 1)$, and a different 1-ladder which has the box $(2, 3)$ in λ and the boxes $(4, 2)$ and $(6, 1)$ not in λ . In the picture below, lines are drawn through the different 1-ladders.




5.2. The ladder crystal. We will construct a new crystal $B(\Lambda_0)^L$ recursively as follows. First, the empty partition $\emptyset$ is the unique highest weight node of our crystal. From $\emptyset$, we will build the crystal by applying the operators $\widehat{f}_i$ for $0 \leq i < \ell$. We define $\widehat{f}_i$ to act on partitions, taking a partition of n to a partition of $n+1$ (or 0) in the following manner. Given $\lambda \vdash n$, first draw all of the i -ladders of λ onto its Young diagram. Label any addable i -box with a $+$, and any removable i -box with a $-$. Now, write down the word of $+$'s and $-$'s by reading from leftmost i -ladder to rightmost i -ladder and reading from top to bottom on each ladder. This is called the *ladder i -signature* of λ . From here, cancel any adjacent $-+$ pairs in the word, until you obtain a word of the form $+\cdots+ -\cdots-$. This is called the *reduced ladder i -signature* of λ . All boxes associated to a $-$ in the reduced ladder i -signature are called *ladder normal i -boxes* and all boxes associated to a $+$ in the reduced ladder i -signature are called *ladder conormal i -boxes*. The box associated to the leftmost $-$ is called the *ladder good i -box* and the box associated to the rightmost $+$ is called the *ladder cogood i -box*. Then we define $\widehat{f}_i\lambda$ to be the partition λ union the ladder cogood i -box. If no such box exists, then $\widehat{f}_i\lambda = 0$. Similarly, $\widehat{e}_i\lambda$ is the partition λ with the ladder good i -box removed. If no such box exists, then $\widehat{e}_i\lambda = 0$. We then define $\widehat{\varphi}_i(\lambda)$ to be the number of ladder conormal i -boxes of λ and $\widehat{\varepsilon}_i(\lambda)$ to be the number of ladder normal i -boxes. It is then obvious that $\widehat{\varphi}_i(\lambda) = \max\{k : \widehat{f}_i^k\lambda \neq 0\}$ and that $\widehat{\varepsilon}_i(\lambda) = \max\{k : \widehat{e}_i^k\lambda \neq 0\}$. For the rest of the paper, $\widehat{\varphi} = \widehat{\varphi}_i(\lambda)$ and $\widehat{\varepsilon} = \widehat{\varepsilon}_i(\lambda)$.

Remark 5.2.1. It remains to be shown that this directed graph is a crystal. To see this, we will show that it is isomorphic to $B(\Lambda_0)$.

Remark 5.2.2. Those who study crystals know that they also have a weight function assigned to the nodes of the graph. In here, just as in the classical description $B(\Lambda_0)$, the weight of a partition λ is given by $wt(\lambda) = \Lambda_0 - \sum_i a_i \alpha_i$, where a_i denotes the number of boxes of λ of residue i .

Example 5.2.3. Let $\lambda = (5, 3, 1, 1, 1, 1)$ and $\ell = 3$. Then there are four addable 2-boxes for λ . In the leftmost 2-ladder (containing box (2,1)) there are no addable (or removable) 2-boxes. In the next 2-ladder (containing box (1,3)) there is an addable 2-box in box (3,2). In the next 2-ladder (containing box (2,4)), there

are two addable 2-boxes, in boxes $(2,4)$ and $(8,1)$. In the last drawn 2-ladder (containing box $(1,6)$) there is one addable 2-box, in box $(1,6)$. There are no removable 2-boxes in λ . Therefore the ladder 2-signature (and hence reduced ladder 2-signature) of λ is $+(3,2) + (2,4) + (8,1) + (1,6)$ (Here, we have included subscripts on the $+$ signs so that the reader can see the correct order of the $+$'s). Hence $\widehat{f}_2\lambda = (6, 3, 1, 1, 1, 1, 1)$, $(\widehat{f}_2)^2\lambda = (6, 3, 1, 1, 1, 1, 1, 1)$, $(\widehat{f}_2)^3\lambda = (6, 4, 1, 1, 1, 1, 1, 1)$ and $(\widehat{f}_2)^4\lambda = (6, 4, 2, 1, 1, 1, 1, 1)$. $(\widehat{f}_2)^5\lambda = 0$. See Figure 5.2 for the 2-string of λ .



From this description, it is not obvious that this is a crystal. However, we will soon show that it is isomorphic to $B(\Lambda_0)$.

Remark 5.2.4. We note that this crystal rule is well defined for all partitions. We will mainly be studying the connected component of $\emptyset$, which is what we will show is isomorphic to $B(\Lambda_0)$.

Remark 5.2.5. For the crystal expert: The weight function of this crystal is exactly the same as the weight function for $B(\Lambda_0)^L$. Explicitly, the weight of λ is $\Lambda_0 - \sum c_i \alpha_i$ where c_i is the number of boxes of λ with residue i . Throughout this paper we will suppress the weight function as it is irrelevant to the combinatorics involved.

We end this section by proving a simple property of $\widehat{e}_i$ and $\widehat{f}_i$.

Lemma 5.2.6. $\widehat{e}_i\lambda = \mu$ if and only if $\widehat{f}_i\mu = \lambda$.

Proof. In the reduced i -signature of μ , the rightmost $+$ will be changed by $\widehat{f}_i$ to a $-$. The resulting word will have no additional cancelation. Therefore, the reduced i -signature of $\lambda = \widehat{f}_i\mu$ will have its leftmost $-$ corresponding to the same box. Thus $\widehat{e}_i\lambda = \mu$. The other direction is similar. $\square$




FIGURE 2. The 2-string of the partition $(5, 3, 1, 1, 1, 1, 1)$ in $B(\Lambda_0)$ for $\ell = 3$ from Example 5.2.3

6. REGULARIZATION

6.1. The operation of regularization. In this section we describe a map from the set of partitions to the set of ℓ -regular partitions. The map is called regularization and was first defined by James (see [8]). For a given λ , move all of the boxes up to the top of their respective ladders. The result is a partition, and that partition is called the *regularization* of λ , and is denoted $\mathcal{R}_\ell \lambda$. Although $\mathcal{R}_\ell$ depends on ℓ , we will usually just write $\mathcal{R}$. The following theorem contains facts about regularization originally due to James [8] (see also [11]).

Theorem 6.1.1. *Let λ be a partition. Then*

- $\mathcal{R}\lambda$ is ℓ -regular
- $\mathcal{R}\lambda = \lambda$ if and only if λ is ℓ -regular.
- If λ is ℓ -regular and $D^\lambda \cong S^\nu$ for some partition ν , then $\mathcal{R}\nu = \lambda$.

Regularization provides us with an equivalence relation on the set of partitions. Specifically, we say $\lambda \sim \mu$ if $\mathcal{R}\lambda = \mathcal{R}\mu$. The equivalence classes are called





FIGURE 3. The first 6 levels of $B(\Lambda_0)^L$ for $\ell = 3$

regularization classes, and the class of a partition λ is denoted $\mathcal{RC}(\lambda) := \{\mu \in \mathcal{P} : \mathcal{R}\mu = \mathcal{R}\lambda\}$.

Example 6.1.2. Let $\lambda = (2, 2, 2, 1, 1, 1)$ and let $\ell = 3$. Then $\mathcal{R}\lambda = (3, 3, 2, 1)$. Also,

$$\mathcal{RC}(\lambda) = \{(2, 2, 2, 1, 1, 1), (2, 2, 2, 2, 1), (3, 2, 1, 1, 1, 1), \\ (3, 2, 2, 2), (3, 3, 1, 1, 1), (3, 3, 2, 1)\}$$



7. DEREGULARIZATION

The goal of this section is to provide an algorithm for finding the smallest partition in dominance order in a given regularization class. It is nontrivial to show that a smallest partition exists. We use this result to show that our new description of the crystal $B(\Lambda_0)^L$ has nodes which are smallest in dominance order in their regularization class. All of the work of this section is inspired by Brant Jones of UC Davis, who gave the first definition of a locked box.

7.1. Locked Boxes. We recall a partial ordering on the set of partitions of n . For two partitions λ and μ of n , we say that $\lambda \leq \mu$ if $\sum_{j=1}^i \lambda_j \leq \sum_{j=1}^i \mu_j$ for all i . This order is usually called the *dominance order*.

Definition 7.1.1. The k^{th} ladder of λ is the ladder which contains the box $(k, 1)$ (this ladder necessarily has residue $-k + 1 \pmod{\ell}$).

Finding all of the partitions which belong to a regularization class is not easy. The definition of locked boxes below formalizes the concept that some boxes in a partition cannot be moved down their ladders if one requires that the new diagram remain a partition.

Definition 7.1.2. For a partition λ , we label boxes of λ as *locked* by the following procedure:

- (1) If a box x has a locked box directly above it (or is on the first row) and every unoccupied space in the same ladder as x , lying below x , has an unoccupied space directly above it then x is locked. Boxes locked for this reason are called type I locked boxes.
- (2) If a box y is locked, then every box to the left of y in the same row is also locked. Boxes locked for this reason are called type II locked boxes.

Boxes which are not locked are called *unlocked*.

Remark 7.1.3. Locked boxes can be both type I and type II.

Example 7.1.4. Let $\ell = 3$ and let $\lambda = (6, 5, 4, 3, 1, 1)$. Then labeling the locked boxes for λ with an L and the unlocked boxes with a U yields the picture below.

L	L	L	U	U	U
L	L	L	U	U	
L	L	U	U		
L	L	U			
L					
L					

The following lemmas follow from the definition of locked boxes.

Lemma 7.1.5. *If (a, b) is locked and $a > 1$ then $(a - 1, b)$ is locked. Equivalently, all boxes which sit below an unlocked box in the same column are unlocked.*

Proof. If (a, b) is a type I locked box then by definition $(a - 1, b)$ is locked. If (a, b) is a type II locked box and not type I then there exists a c so that (a, c) is a type I locked box with $c > b$. But then by definition of type I locked box, $(a - 1, c)$ is locked. Then $(a - 1, b)$ is a type II locked box. $\square$

Lemma 7.1.6. *If there is a locked box in space (α, β) and there is a box in space $(\alpha - (\ell - 1), \beta + 1)$ then the box $(\alpha - (\ell - 1), \beta + 1)$ is locked.*

Proof. To show this, suppose that $(\alpha - (\ell - 1), \beta + 1)$ is unlocked. Then let $(\gamma, \beta + 1)$ be the highest unlocked box in column $\beta + 1$. This is unlocked because it violates the type I locked condition. Then the box $(\gamma + \ell - 1, \beta)$ violates the type I locked condition, and it will not have a locked box directly to the right of it ($(\gamma, \beta + 1)$ is unlocked, so $(\gamma + \ell - 1, \beta + 1)$ will be unlocked if it is occupied, by Lemma 7.1.5). Hence $(\gamma + \ell - 1, \beta)$ is unlocked, so (α, β) must be unlocked since it sits below $(\gamma + \ell - 1, \beta)$ (by Lemma 7.1.5), a contradiction. $\square$

For two partitions λ and μ in the same regularization class, there are many ways to move the boxes in λ on their ladders to obtain μ . We define an *arrangement of μ from λ* to be an assignment of each box in the Young diagram of λ to a box in the Young diagram of μ . An arrangement will be denoted by a set of ordered pairs (x, y) with $x \in \lambda$ and $y \in \mu$, where both x and y are in the same ladder, and each $x \in \lambda$ and $y \in \mu$ is used exactly once. Such a pair (x, y) denotes that the box x from λ is moved into position y in μ . x is said to *move up* if the position y which it is paired with is higher in the ladder than x is. Similarly, x is said to *move down* if

the position y is lower in the ladder than x is. x is said to *stay put* if the position y in μ has the same coordinates as position x in λ .

Remark 7.1.7. We introduce the notation $\mathcal{B}(p)$ to denote the position paired with p in an arrangement $\mathcal{B}$, i.e. $(p, \mathcal{B}(p)) \in \mathcal{B}$.

Example 7.1.8. $\lambda = (3, 3, 1, 1, 1)$ and $\mu = (2, 2, 2, 2, 1)$ are in the same regularization class when $\ell = 3$ (see Example 6.1.2). One possible arrangement of μ from λ would be $\mathcal{B} = \{((1, 1), (1, 1)), ((1, 2), (1, 2)), ((1, 3), (5, 1)), ((2, 1), (2, 1)), ((2, 2), (2, 2)), ((2, 3), (4, 2)), ((3, 1), (3, 1)), ((4, 1), (4, 1)), ((5, 1), (3, 2))\}$. This corresponds to moving the labeled boxes from λ to μ in the corresponding picture below. Note that $(5, 1)$ moves up, $(1, 3)$ and $(2, 3)$ move down, and all other boxes stay put.

1	2	3	1	2
4	5	6	4	5
7			7	9
8			8	6
9			3	

7.2. Algorithm for finding the smallest partition in a regularization class.

For any partition λ , to find the smallest partition (with respect to dominance order) in a regularization class we first label each box of λ as either locked or unlocked as above. Then we create an arrangement $\mathcal{S}_\lambda$ which moves all unlocked boxes down their ladders, while keeping these unlocked boxes in order (from bottom to top), while locked boxes do not move. The partition obtained from following the arrangement $\mathcal{S}_\lambda$ will be denoted $\mathcal{S}\lambda$. It is unclear that this algorithm will yield the smallest partition in $\mathcal{RC}(\lambda)$, or even that $\mathcal{S}\lambda$ is a partition. In this subsection, we resolve these issues.

Proposition 7.2.1. *Let λ and μ be partitions in the same regularization class. Then there exists an arrangement of μ from λ such that all of the boxes in λ which are locked do not move down.*

Proof. To find a contradiction, we suppose that any arrangement of μ from λ must have at least one locked box in λ which moves down.

For any arrangement $\mathcal{C}$, we let $x_{\mathcal{C}}$ denote the highest rightmost locked box in λ which moves down in $\mathcal{C}$. Since all arrangements of μ from λ have at least one locked box in λ which moves down, $x_{\mathcal{C}}$ is defined for all $\mathcal{C}$.

Among all arrangements of μ from λ , let $\mathcal{D}$ be one which has $x_{\mathcal{D}}$ in the lowest, leftmost position. Let $x = x_{\mathcal{D}} = (a, b)$.

We will exhibit a box w on the same ladder as x , such that $\mathcal{D}(w)$ is equal to or above (a, b) in μ and either w is below x or w is above x and unlocked. If such a box exists, then letting $\mathcal{A} = (\mathcal{D} \setminus \{(x, \mathcal{D}(x)), (w, \mathcal{D}(w))\}) \cup \{(x, \mathcal{D}(w)), (w, \mathcal{D}(x))\}$ will yield a contradiction, as $x_{\mathcal{A}}$ will be in a position to the left of and/or below x , which contradicts our choice of $\mathcal{D}$.

If there exists a w in the same ladder as x , below x , with $\mathcal{D}(w)$ at or above (a, b) in μ then we are done. So now we assume that every box below x in the ladder for x does not move on or above x .

There are two cases to consider:

Case I: x is a type II locked box and not a type I locked box.

In this case, there is a locked box directly to the right of x (definition of a type II lock), which we label y .

If y stays put according to $\mathcal{D}$, (i.e. $(y, y) \in \mathcal{D}$) then some box z in the same ladder as x in λ must move into the position for x according to $\mathcal{D}$, i.e. $(z, x) \in \mathcal{D}$. Let $w = z$. Since w will move into position x , w is above x , so it must be unlocked (because $x = x_{\mathcal{D}}$ was the highest locked box which moved down according to $\mathcal{D}$) and we are done.

If y moves according to $\mathcal{D}$ then it must move up (since it is to the right of x and locked).

If there are any boxes $\tilde{x}$ in the ladder of x , above x , which are at the end of their row, then $\tilde{x}$ is unlocked (it is not a type II lock because there are no boxes to the right of it, and it can't be a type I lock because x is not a type I lock). If $\tilde{x}$ were to move according to $\mathcal{D}$ to some box on or above x , then we could use $w = \tilde{x}$ and be done. So otherwise we assume all such boxes move down below x according to $\mathcal{D}$.

Similarly, if there are any boxes $\hat{x}$ above x , in the ladder for x , which are directly to the left of a box $\hat{y}$ in the ladder for y , and the box $\hat{y}$ moves down according to $\mathcal{D}$, then the box $\hat{x}$ must be unlocked (its not a type II lock because $\hat{y}$ is unlocked,

being above x , and its not a type I lock because x is not). If $\hat{x}$ moves to a position on or above x , then we could use $w = \hat{x}$ and be done. Otherwise, we must assume that all such $\hat{x}$ move down below x .

Assuming we cannot find any w by these methods, we let

- k denote the number of boxes on the ladder of x , above (a, b) , in λ ,
- j denote the number of boxes on the ladder of y , above (a, b) , in λ ,
- k' denote the number of boxes on the ladder of x , above (a, b) , in μ ,
- j' denote the number of boxes on the ladder of y , above (a, b) , in μ ,
- m denote the number of boxes on the ladder of y , above (a, b) , in λ which move down (i.e. the number of boxes which are of the form $\hat{y}$ above).

The number of boxes of the form $\tilde{x}$ is then $k - j$. Also, $j' \geq j - m + 1$, since the m boxes need not move below position (a, b) , but the box in position y moves above position (a, b) . $k' \geq k - m - (k - j) = j - m$ since the number of boxes in ladder x will go down by at least m for the boxes of the form $\hat{x}$ and $k - j$ for the boxes of the form $\tilde{x}$ (by assumption boxes in the ladder for x cannot move up from below). Hence $k' \leq j - m < j - m + 1 \leq j'$. This is a contradiction, as there must be at least as many boxes above x on the ladder of x as there are on the ladder of y for μ to be a partition.

Case II: x is a type I locked box.

We will show that this case cannot occur. Specifically we will show that the assumption leads to a contradiction of λ being a partition.

Since x is a type I lock, the number of boxes on the ladder for x on or below row a in λ must be strictly greater than the number of boxes on the ladder of $(a - 1, b)$ on or below row a . Since x is moving down, some box in the ladder of $(a - 1, b)$ above row a must move down (since no box on the ladder for x , below x , moves above x , by assumption). Since x is the highest locked box which can move down, this must be an unlocked box. Hence there is an unlocked box in the ladder of $(a - 1, b)$ which is in a row above x .

The existence of an unlocked box in the ladder for $(a - 1, b)$ above row a implies that there is an empty position on the ladder for $(a - 1, b)$ somewhere above row a . This is implied by Lemma 7.1.6 above, since if every space on the ladder $(a - 1, b) \in \lambda$ above row a was occupied, then all of those boxes would be locked.

Let m be the column which contains the lowest such empty position. The coordinates for such a position would then be $(a - 1 - (m - b)(\ell - 1), m)$. Let (c_1, d_1) be the highest empty position on the ladder for x below x (we know a space must exist since x is moved down). Since x is a type I lock, $(c_1 - 1, d_1)$ is also an empty position. We know that $(a - 1, b)$ is locked since x is a type I lock. If $(a - 1, b)$ is also a type I lock then the space $(c_1 - 2, d_1)$ must also be empty. Continuing this, if $(a - (\ell - 2), b)$ were a type I lock then $(c_1 - (\ell - 1), d_1)$ would have to be empty, but this would contradict the fact that there should be a box in $(c_1 - (\ell - 1), d_1 + 1)$ since it is in the ladder for x and (c_1, d_1) was chosen to be the the highest empty position on the ladder for x below x .

So there exists a k_1 so that $(a - k_1, b)$ is a type II lock, with $k_1 \leq \ell - 2$. Since $(a - k_1, b)$ is a type II lock, there exists a type I locked box in a position $(a - k_1, n_1)$ with $b < n_1 < m$. Let (c_2, d_2) denote the highest empty position in the ladder for $(a - k_1, n_1)$ below $(a - k_1, n_1)$ ((c_2, d_2) exists because the position in column d_1 on the ladder for $(a - k_1, n_1)$ is empty). Since $(a - k_1, n_1)$ is a type I lock, the box $(c_2 - 1, d_2)$ is also empty. Continuing as above, if $(a - k_1 - (\ell - 2), n_1)$ were a type I lock, then $(c_2 - (\ell - 1), d_2)$ would be empty, which contradicts our choice of (c_2, d_2) . Hence there exists a $k_2 \leq \ell - 2$ so that $(a - k_1 - k_2, n_1)$ is a type II lock. This implies that there is a type I lock in some position $(a - k_1 - k_2, n_2)$ with $n_1 < n_2 < m$. Continuing this, we get a sequences for k and n with each $k_i \leq \ell - 2$ and $b < n_1 < n_2 < \dots < n_i < m$.

I claim that each of the boxes $(a - \sum_i k_i, n_i)$ are in a ladder below the ladder of $(a - 1, b)$. If this is the case then the sequences for k and n would eventually have to produce a type I locked box in column m or greater, below row $a - 1 - (m - b)(\ell - 1)$. This contradicts that λ is a partition, since there is no box in position $(a - 1 - (m - b)(\ell - 1), m)$.

To show the claim, we just note that each successive type I locked box comes from moving up at most $\ell - 2$ and to the right at least one space. These are clearly in a ladder below the ladder of $(a - 1, b)$, since ladders move up $\ell - 1$ boxes each time they move one box to the right.

□

Corollary 7.2.2. *Fix $n \in \mathbb{N}$. Suppose for any partition $\lambda \vdash n$, $\mathcal{S}\lambda$ is also a partition. Then $\mathcal{S}\lambda$ is the smallest partition (in dominance order) in the regularization class of λ . Furthermore, all boxes of $\mathcal{S}\lambda$ are locked.*

Proof. Let μ be any partition in the regularization class of λ . Then by Proposition 7.2.1 above, we can choose an arrangement $\mathcal{B}$ so that all of the boxes which we must move down from λ to form μ are unlocked boxes. But the arrangement $\mathcal{S}\lambda$ moves all unlocked boxes as far down as they can go, so $\mathcal{S}\lambda \leq \mu$. For the second statement, assume some box of $\mathcal{S}\lambda$ is unlocked. Then $\mathcal{S}_{\mathcal{S}\lambda}$ must move some boxes down. Following the arrangement $\mathcal{S}_{\mathcal{S}\lambda}$ yields another partition $\mathcal{S}^2\lambda$ which is smaller than $\mathcal{S}\lambda$, which contradicts the first statement²³ of this corollary. $\square$

Proposition 7.2.3. *Let λ be a partition. Then $\mathcal{S}\lambda$ is a partition.*

Proof. There are two possibilities we must rule out: That an unlocked box in λ which was moved down via the arrangement $\mathcal{S}\lambda$ has an empty position directly to the left, or that it has an empty position directly above. If we show that the algorithm never leaves an empty position above a moved unlocked box, then as a consequence we can easily prove that we get no empty positions to the left of a moved box.

Suppose the algorithm never leaves an empty position above a moved box. If there was an empty position x directly to the left of a moved box $\mathcal{S}\lambda(y)$ then below x there would be a box z on the same ladder as $\mathcal{S}\lambda(y)$. If z was not moved, then it must have been locked, which implies all of the boxes above it were locked, including the box which was in position x , contradicting that position x is empty. Otherwise, z moved, so our assumption implies that z has no empty position above it. Applying this procedure again, we can determine that there must be a box directly above the box directly above z . Applying this procedure will eventually imply that there must be a box in position x . Therefore our goal is to show that moving down all unlocked boxes produces no box below an empty position.

We prove this by induction on n , the number of boxes in λ . The $n = 1$ case is clear. We assume that if η is a partition of $k < n$ then $\mathcal{S}\eta$ has no box below an empty position (and hence is a partition by the previous paragraph). We let λ be any partition of n and we will show that $\mathcal{S}\lambda$ has no box below an empty position.

The inductive proof is broken into three cases.

The first case is that there exists an i so that $\lambda_1 = \lambda_i$ and the box $x = (i, \lambda_i)$ is locked. Then all of the boxes in the first i rows are also locked. Let $\mu = \lambda \setminus \{\text{first } i \text{ rows}\}$. Boxes are locked in μ if and only if they are locked in $\mu \subset \lambda$. Since $|\mu| < |\lambda|$, $\mathcal{S}\mu$ has no box below an empty position. We append the first i rows back on top of $\mathcal{S}\mu$ to form $\mathcal{S}\lambda$. Hence $\mathcal{S}\lambda$ has no box below an empty position.

The second case is when there does not exist such an i . Let j be so that $\lambda_1 = \lambda_j \neq \lambda_{j+1}$. If $j > 1$ then we let x be the box (j, λ_j) and let $y = (j-1, \lambda_{j-1})$. Let $\mu = \lambda \setminus \{x\}$. Let $\nu = \lambda \setminus \{x, y\}$. Note that boxes in μ (and ν) are locked if and only if they are locked in λ . Since $|\mu| < |\lambda|$, we can form a partition $\mathcal{S}\mu$ by bringing down all unlocked boxes. Now we place x into the lowest empty position on the ladder of x in $\mathcal{S}\mu$. If it is not a partition, it is because there is an empty position above where x was placed. If this is the case, then the box y was moved down below where x was pushed down, and y has a box z below it (if it didn't then x would move below y). But then in ν when we move down all of the boxes we would have z below an empty position (where y is in μ). But $|\nu| < |\lambda|$, so $\mathcal{S}\nu$ should not have any empty positions.

Lastly, if $j = 1$ then we let $x = (1, \lambda_1)$. Since x is unlocked, there is at least one empty position in the ladder for x which has a box above it. Let $\mu = \lambda \setminus \{x\}$. Boxes are locked in μ if and only if they were locked in λ . We move all the unlocked boxes of μ down; by induction and Corollary 7.2.2 this is the smallest partition in the regularization class of μ , denoted $\mathcal{S}\mu$. Since the number of boxes on the ladder for x was at most the number of boxes on the ladder directly above x in λ (this is because x cannot be a type I lock), the number of boxes on the ladder for x is strictly less than the number on the ladder above x in μ . Therefore, there exists an empty position on the ladder for x directly below a box in $\mathcal{S}\mu$. We let (a, b) be the lowest such empty position. If (a, b) is the lowest empty position in $\mathcal{S}\mu$ on the ladder for x , then moving x into that space will yield a partition which is obtained from moving x into the lowest empty position in its ladder. If not, then there must be an empty position on the ladder for x directly below an empty position, all below row a . Let (c, d) be the highest such empty position on the ladder for x below (a, b) . Let (m_0, d) be the lowest box in column d (m_0 is at least $c - \ell$, since

there is a box in the space $(c - (\ell - 1), d + 1)$). Let (m_1, b) be the box in column b in the same ladder as $(m_0 + 1, d)$. By Corollary 7.2.2, all of the boxes of $\mathcal{S}\mu$ are locked. But (m_1, b) has the space $(m_0 + 1, d)$ in the same ladder, so if (m_1, b) is locked it must be because a box to the right of it is locked (i.e. it is a type II locked box). Call this box (m_1, b_1) . We continue by letting (m_2, b_1) be the box in the same ladder as $(m_0 + 1, d)$ in column b_1 . Similarly, since this box is locked, there must be a box to the right of it which is locked. That box will be (m_2, b_2) . This process must eventually conclude at step (m_k, b_{k-1}) where b_{k-1} is less than the column of x (since every box in μ has a column value less than that of x) and the space $(m_k, b_{k-1} + 1)$ is empty. We just need to show that the box (m_k, b_{k-1}) is actually a box in the diagram of λ (i.e. that m_k is at least 1). In fact, $(m_0 + 1, d)$ is either in the ladder directly to the left of x or below this ladder. Hence (m_k, b_{k-1}) is either in the ladder left of x or below. It must also be in a column strictly to the left of x . Therefore (m_k, b_{k-1}) is a box in λ . But then the box (m_k, b_{k-1}) must be unlocked, since it has no locked boxes to the right and the empty position $(m_0 + 1, d)$ in the same ladder below it. This contradicts all of the boxes of $\mathcal{S}\mu$ being locked. $\square$

Example 7.2.4. Continuing from the example above ($\lambda = (6, 5, 4, 3, 1, 1)$ and $\ell = 3$), we move all of the unlocked boxes down to obtain the smallest partition in $\mathcal{RC}(\lambda)$, which is $\mathcal{S}\lambda = (3, 3, 2, 2, 2, 2, 1, 1, 1, 1)$. The boxes labeled L are the ones which were locked in $(6, 5, 4, 3, 1, 1)$ (and did not move).

L	L	L
L	L	L
L	L	
L	L	
L		
L		

Theorem 7.2.5. *$\mathcal{S}\lambda$ is the unique smallest partition in its regularization class with respect to dominance order. It can be classified as being the unique partition (in its regularization class) which has all locked boxes.*

Proof. This follows from Corollary 7.2.2 and Proposition 7.2.3. $\square$

7.3. The nodes of $B(\Lambda_0)^L$ are smallest in dominance order. The nodes of $B(\Lambda_0)^L$ have been defined recursively by applying the operators $\widehat{f}_i$. We now give a simple description which determines when a partition is a node of $B(\Lambda_0)^L$.

Proposition 7.3.1. *Let λ be a partition of n . Let $\mathcal{RC}(\lambda)$ be its regularization class. If λ is a node of $B(\Lambda_0)^L$ then λ is the smallest partition in $\mathcal{RC}(\lambda)$ with respect to dominance order.*

Proof. The proof is by induction on the number of boxes in λ . Suppose we have a partition in $B(\Lambda_0)^L$ which is smallest in its regularization class. Equivalently, all of the boxes of λ are locked. We want to show that $\widehat{f}_i\lambda$ is still smallest in its regularization class. Let $x = \widehat{f}_i\lambda \setminus \lambda$, so that $\widehat{f}_i\lambda$ is just λ with the addable i -box x inserted. There are two cases to consider.

The first is that the insertion of x into λ makes x into an unlocked box. Since the position y above x must be locked (or x is in the first row), there must exist an empty position x' in the same ladder as x which has a box y' directly above x' in the ladder of y . Also, the box z to the left of x must be a type I lock, since x was not in λ (implying z could not be type II). But then the position z' to the left of x' must have a box, since there is a box in the position above z' (the box to the left of y'). Since λ was smallest in dominance order, it could not have had a $+$ below a $-$ on the same ladder (otherwise the $-$ could be moved down to the $+$, contradicting that λ was smallest in dominance order). Hence there is no $-$ between the two $+$'s in positions x and x' of λ . This means that the crystal rule would have chosen to add to x' instead of x , a contradiction.

The second case is that the insertion of x into λ unlocks a box. If this is the case then let y be the position above x . The only case to consider is adding the box x unlocks some box above the row of x , on the ladder directly below x . Let z denote the lowest such box. If there is a box x' directly above z and no box directly to the

right of x' then x' would have been unlocked in λ , since the box y is in λ but x is not. This contradicts the assumption that λ was the smallest.

The only other possibility is that the box x' above z has a box directly to the right of it (this still works even if z is in the first row of the partition). If there was a box to the right of z , then that box would be locked since it was locked in λ . But then z would be locked independently of the addition of x . So z is at the end of its row. The space one box to the right of z (let us name it w) must therefore be empty and have the same residue as x . It is an addable box, so it will contribute a $+$ to the i -ladder-signature. There are no $-$ boxes on the ladder for x below x , because if there was a removable i -box on the same ladder as x below x then the box z would have been unlocked in λ . Therefore, no $-$ will cancel the $+$ from space w . Since w is on the ladder past the ladder for x , this yields a contradiction as $\widehat{f}_i$ should have added a box to position w .

□

One can view $B(\Lambda_0)$ as having nodes $\{\mathcal{RC}(\lambda) : \lambda \vdash n, n \geq 0\}$. The usual model of $B(\Lambda_0)$ takes the representative $\mathcal{R}\lambda \in \mathcal{RC}(\lambda)$, which happens to be the largest in dominance order. Here, we will take a different representative of $\mathcal{RC}(\lambda)$, the partitions $\mathcal{S}\lambda$, which are smallest in dominance order. One must then give a description of the edges of the crystal graph. We will show in Section 10 that the crystal operators $\widehat{f}_i$ and $\widehat{e}_i$ are the correct operators to generate the edges of the graph. In other words, we will show that the crystal $B(\Lambda_0)^L$ constructed above is indeed isomorphic to $B(\Lambda_0)$.

8. EXTENDING THEOREMS TO THE CRYSTAL $B(\Lambda_0)^L$

8.1. Motivation. In this section, we prove analogues of Theorems 4.3.1 and 4.3.2 for our newly defined crystal $B(\Lambda_0)^L$, with $(\ell, 0)$ -JM partitions taking the place of ℓ -partitions. We start with a few lemmas about $B(\Lambda_0)^L$.

8.2. Crystal theoretic results for $B(\Lambda_0)^L$ and $(\ell, 0)$ -JM partitions. Our goal will be to show that all $(\ell, 0)$ -JM partitions are in $B(\Lambda_0)^L$. First we prove that all ℓ -cores are in $B(\Lambda_0)^L$.

Lemma 8.2.1. *If λ is an ℓ -core, then $\widehat{\varphi} = \varphi$ and $\widehat{f}_i^\varphi \lambda = \tilde{f}_i^\varphi \lambda$. Hence all ℓ -cores are nodes of $B(\Lambda_0)^L$.*

Proof. ℓ -cores have either all addable or all removable i -boxes for any fixed i . Therefore, for a core λ we have that $\varphi = \widehat{\varphi}$. Thus $\widehat{f}_i^\varphi$ and $\widetilde{f}_i^\varphi$ will both add all addable i -boxes to λ , just in different relative orders.

For the last statement, recall that applying the crystal operators $\widetilde{f}_i^\varphi$ will generate all ℓ -cores (starting with the empty partition $\emptyset$). $\square$

The following lemma is a well known recharacterization of ℓ -cores.

Lemma 8.2.2. *A box x has a hook length divisible by ℓ if and only if there exists a residue i so that the last box in the row of x has residue i and the last box in the column of x has residue $i + 1$. In particular, any partition which has such a box x is not an ℓ -core.*

The upcoming lemma is used only in the proof of Lemma 8.2.4.

Lemma 8.2.3. *If λ is an $(\ell, 0)$ -JM partition then for every $0 \leq i < \ell$, there is no i -ladder in the Young diagram of λ which has a $-$ above a $+$.*

Proof. Suppose λ is an $(\ell, 0)$ -JM partition. Then $\lambda \approx (\mu, r, s, \rho, \sigma)$. First suppose that $\mu = \emptyset$. Let j be the residue of the addable box in the first row of λ . Then all removable boxes have residue either $j - 1$ (in the first r rows) or $j + 1$ (in the first s columns), so no ladder contains both a $-$ and a $+$.

Suppose $\mu \neq \emptyset$. The only possible way of having a $-$ above a $+$ in a ladder is to have a removable box of residue j on or above row $r + 1$ and an addable box of residue j on or to the left of column $s + 1$. But then the core μ has a box of residue j at the end of its first row (row $r + 1$ of λ) and a box of residue $j + 1$ at the end of its first column (on column $s + 1$ of λ). This implies that μ isn't a core, by Lemma 8.2.2, since $\ell \mid h_{(1,1)}^\mu$. $\square$

The following Lemma 8.2.4 will be used in this section for proving our crystal theorem generalizations for $(\ell, 0)$ -JM partitions.

Lemma 8.2.4. *Let λ be an $(\ell, 0)$ -JM partition. Then its ladder i -signature is the same as the reduced ladder i -signature. In other words, there is no $-+$ cancelation in the ladder i -signature of λ .*

Proof. Suppose there is a $-+$ cancelation in the ladder i -signature of an $(\ell, 0)$ -JM partition λ . By Lemma 8.2.3, it must be that a removable i -box occurs on a ladder

to the left of a ladder which contains an addable i -box. Suppose the removable i -box is in box (a, b) and the addable i -box is in box (c, d) . We will suppose that $a > c$ (the case $a < c$ is similar). Then $\ell \mid h_{(c,b)}^\lambda$. Also $h_{(a,b)}^\lambda = 1$ since (a, b) is a removable box. If all of $h_{(c,k)}^\lambda$ are divisible by ℓ for $k \in \{b+1, b+2, \dots, d-1\}$, then the box (a, b) could not be in a ladder to the left of the ladder of (c, d) . Therefore there is a $b < j < d$ so that $\ell \nmid h_{(c,j)}^\lambda$. Hence λ is not an $(\ell, 0)$ -JM partition. $\square$

8.3. Generalizations of the crystal theorems to $B(\Lambda_0)^L$. We will now prove analogues of Theorems 4.3.1 and 4.3.2 for $(\ell, 0)$ -JM partitions in the ladder crystal $B(\Lambda_0)^L$.

Theorem 8.3.1. *Suppose that λ is an $(\ell, 0)$ -JM partition and $0 \leq i < \ell$. Then*

- (1) $\widehat{f}_i^{\widehat{\varphi}} \lambda$ is an $(\ell, 0)$ -JM partition,
- (2) $\widehat{e}_i^{\widehat{\varphi}} \lambda$ is an $(\ell, 0)$ -JM partition.

Proof. We will prove (1); (2) follows similarly. Suppose that $\lambda \approx (\mu, r, s, \rho, \sigma)$ has an addable m -box in the first row, and an addable n -box in the first column for two residues m, n . If $m \neq i \neq n$ then $\widehat{f}_i^{\widehat{\varphi}}$ will only add boxes to the core μ in the Young diagram of λ . But $\widehat{f}_{i-r+s}^{\widehat{\varphi}} \mu$ will again be a core, by Lemma 8.2.1. Hence $\widehat{f}_i^{\widehat{\varphi}} \lambda \approx (\widehat{f}_{i-r+s}^{\widehat{\varphi}} \mu, r, s, \rho, \sigma)$.

If $m = i \neq n$ then the partition $\nu \approx (\mu, r, 0, \rho, \emptyset)$ is an ℓ -partition. From Lemma 8.2.4, we have no cancelation of $-+$ in λ , so that $\widehat{f}_{i-s}^{\widehat{\varphi}^{i-s}(\nu)} \nu = \tilde{f}_{i-s}^{\varphi^{i-s}(\nu)} \nu$. By Theorem 4.3.1, $\widehat{f}_{i-s}^{\widehat{\varphi}^{i-s}(\nu)} \nu$ is an ℓ -partition. Say $\widehat{f}_{i-s}^{\widehat{\varphi}^{i-s}(\nu)} \nu \approx (\mu', r', 0, \rho', \emptyset)$. Then $\widehat{f}_i^{\widehat{\varphi}} \lambda \approx (\mu', r', s, \rho', \sigma)$. A similar argument works when $m \neq i = n$ by using that the transpose of $(\mu, 0, s, \emptyset, \sigma)$ is an ℓ -partition.

We now suppose that $m = i = n$. In this case, λ has an addable i -box in the first $r+1$ rows and $s+1$ columns. It may also have addable i -boxes within the core μ . λ has no removable i -boxes. Thus we get $\widehat{f}_i^{\widehat{\varphi}} \lambda \approx (\widehat{f}_{i-r+s}^{\widehat{\varphi}^{i-r+s}(\mu)} \mu, r, s, \rho, \sigma)$ is an $(\ell, 0)$ -JM partition. $\square$

Theorem 8.3.2. *Suppose that λ is an $(\ell, 0)$ -JM partition. Then*

- (1) $\widehat{f}_i^k \lambda$ is not an $(\ell, 0)$ -JM partition for $0 < k < \widehat{\varphi} - 1$,
- (2) $\widehat{e}_i^k \lambda$ is not an $(\ell, 0)$ -JM partition for $1 < k < \widehat{\varepsilon}$.

Proof. We will prove (1), and we claim that (2) is similar. We will prove this by showing that there must exist boxes (a, b) , (a, y) and (x, b) in the Young diagram

of λ such that $\ell \mid h_{(a,b)}^\lambda$ and $\ell \nmid h_{(a,y)}^\lambda, \ell \nmid h_{(x,b)}^\lambda$. Since $0 < k < \widehat{\varphi} - 1$ there are at least two ladder conormal i -boxes in $\widehat{f}_i^k \lambda$ and at least one ladder normal i -box. Label the normal box n_1 . Label the conormal boxes n_2 and n_3 (without loss of generality, n_2 is in a row above n_3). We will show that any partition with at least one ladder normal i -box and at least two ladder conormal i -boxes cannot be an $(\ell, 0)$ -JM partition. There are three cases to consider.

The first case is that n_1 is above n_2 and n_3 . Then the hook length in the row of n_1 and column of n_3 is divisible by ℓ , but the hook length in the row of n_2 and column of n_3 is not. Also, the hook length for box n_1 is 1, which is not divisible by ℓ .

The second case is that n_1 is in a row between the row of n_2 and n_3 . In this case, ℓ divides the hook length in the row of n_1 and column of n_3 . Also ℓ does not divide the hook length in the row of n_2 and column of n_3 , and the hook length for the box n_1 is 1.

The last case is that n_1 is below n_2 and n_3 . In this case, ℓ divides the hook length in the column of n_1 and row of n_2 , but ℓ does not divide the hook length in the column of n_3 and row of n_2 . Also the hook length for the box n_1 is 1.

□

8.4. All $(\ell, 0)$ -JM partitions are nodes of $B(\Lambda_0)^L$. This short subsection contains the proof that all $(\ell, 0)$ -JM partitions occur as nodes of $B(\Lambda_0)^L$. This surprising fact is extremely important to the proofs of Theorems 4.3.5 and 4.3.6.

Theorem 8.4.1. *If λ is an $(\ell, 0)$ -JM partition then λ is a node of $B(\Lambda_0)^L$.*

Proof. The proof is by induction on the size of a partition. If the partition has size zero then it is the empty partition which is an $(\ell, 0)$ -JM partition and is a node of the crystal $B(\Lambda_0)^L$.

Suppose $\lambda \vdash n$ is an $(\ell, 0)$ -JM partition. Let i be a residue so that λ has at least one ladder conormal box of residue i . We can find such a box since no $-+$ cancellation exists by Lemma 8.2.4. Define μ to be $\widehat{e}_i^\varepsilon \lambda$. Then $\mu \vdash (n - \varepsilon)$ is an $(\ell, 0)$ -JM partition by Theorem 8.3.1, of smaller size than λ . By induction μ is a node of $B(\Lambda_0)^L$. But $\widehat{f}_i^\varepsilon \mu = \widehat{f}_i^\varepsilon \widehat{e}_i^\varepsilon \lambda = \lambda$ by Lemma 5.2.6, so λ is a node of $B(\Lambda_0)^L$.

□

9. REINTERPRETING THE CRYSTAL RULE OF $B(\Lambda_0)$

In this short section we prove some lemmas necessary for our main theorem (that the crystals $B(\Lambda_0)$ and $B(\Lambda_0)^L$ are isomorphic). We also reinterpret the crystal rule on the classical $B(\Lambda_0)$ in terms of the new crystal rule.

9.1. Two lemmas needed for crystal isomorphism.

Lemma 9.1.1. *If $\lambda \in B(\Lambda_0)^L$ then for every $0 \leq i < \ell$, there is no i -ladder in the Young diagram of λ which has a $-$ above a $+$ (i.e. there do not exist boxes (a, b) and (c, d) on the same i -ladder of λ so that (a, b) is removable, (c, d) is addable and $a < c$).*

Proof. If there was a $-$ box above an empty $+$ box on the same ladder, then one could move that $-$ box down to the $+$ space to form a new partition in the same regularization class. But this cannot happen since the nodes of $B(\Lambda_0)^L$ are smallest in dominance order. $\square$

Lemma 9.1.2. *If $\lambda \in B(\Lambda_0)$ then for every $0 \leq i < \ell$, there is no i -ladder in the Young diagram of λ which has a $+$ above a $-$ (i.e. there do not exist boxes (a, b) and (c, d) on the same i -ladder of λ so that (a, b) is addable, (c, d) is removable and $a < c$).*

Proof. Since λ is in $B(\Lambda_0)$, it is ℓ -regular. Suppose such positions (a, b) and (c, d) exist. Then the rim hook starting at box $(a, b - 1)$ and following the border of λ down to the (c, d) box will cover exactly $(b - d)\ell$ boxes. However, this border runs over only $b - d$ columns, so there must exist a column which contains at least ℓ boxes at the ends of their rows. This contradicts λ being ℓ -regular. $\square$


9.2. Reinterpreting the classical crystal rule. This short theorem proves that the classical crystal rule can be interpreted in terms of ladders. In fact the only difference between the classical rule and the ladder crystal rule is that ladders are read bottom to top instead of top to bottom.

Theorem 9.2.1. *The i -signature (and hence reduced i -signature) of an ℓ -regular partition λ in $B(\Lambda_0)$ can be determined by reading from its leftmost ladder to rightmost ladder, bottom to top.*

Proof. If positions x and y contain $+'s$ (or $- 's$) of the same residue with the ladder of x to the left of the ladder of y , then by the regularity of λ , x will be in a row below y . Hence reading the i -signature up ladders from left to right is equivalent to reading up the rows of λ from bottom to top.

□

Example 9.2.2. $\lambda = (6, 5, 3, 3, 2, 2, 1)$ and $\ell = 3$. Suppose we wanted to find the 2-signature for λ . Then we could read from leftmost ladder (in this case, the leftmost ladder relevant to the 2-signature is the one which contains position $(8, 1)$) to rightmost ladder. Inside each ladder we read from bottom to top. The picture below shows the positions which correspond to addable and removable 2-boxes, with their ladders. The 2-signature is $+ - - -$.



10. REGULARIZATION AND CRYSTAL ISOMORPHISM

The results of this section come from ideas originally sketched out with Steve Pon in the summer of 2007.

10.1. Crystal isomorphism. We will now prove that our crystal $B(\Lambda_0)^L$ from Section 5 is indeed isomorphic to the crystal $B(\Lambda_0)$. We first classify positions in a Young diagram. Let λ be an arbitrary partition and suppose $x = (i, j)$ with $i, j \geq 0$. We use the convention that x is in λ if either of i, j are not positive. We say the type of x with respect to λ is:

- (a) if the box x is in λ and the box $(i + 1, j + 1)$ is in λ .
- (b) if the box x is in λ , the boxes $(i + 1, j)$ and $(i, j + 1)$ are in λ and $(i + 1, j + 1)$ is not in λ .
- (c) if the box x is in λ , $j > 0$, the box $(i + 1, j)$ is in λ and $(i, j + 1)$ is not in λ .

- (d) if the box x is in λ , the box $(i, j + 1)$ is in λ and $(i + 1, j)$ is not in λ .
- (e) if the box x is in λ and $(i + 1, j), (i, j + 1)$ are not in λ .
- (f) if $j = 0$ and the box $(i, 1)$ is not in λ .
- (g) if the box x is not in λ , and $(i - 1, j - 1)$ is of type (e).
- (h) if the box x is not in λ , and $(i - 1, j - 1)$ is of type (c).
- (i) if the box x is not in λ , and $(i - 1, j - 1)$ is of type (d).
- (j) if the box x is not in λ , and $(i - 1, j - 1)$ is of type (b).
- (k) if the box x is not in λ , and $(i - 1, j - 1)$ is of type (f) or not in λ .

Remark 10.1.1. A box x is removable if and only if it is type (e) and addable if and only if it is type (j)

Example 10.1.2. Let $\lambda = (4, 2, 1, 1)$. Then the boxes are labeled by type in the following picture.

	a	a	a	a	b	d	d
a	a	b	d	fe	j	i	
a	b	e	j	i	g	k	
a	c	j	g	k	k	k	
b	e	h	k	k	k	k	
f	j	g	k	k	k	k	
f	k	k	k	k	k	k	

Definition 10.1.3. The k^{th} ladder of λ will refer to all of the boxes (i, j) of λ which are on the same ladder as $(k, 1)$.

Definition 10.1.4. The number of boxes of type α in ladder k of the partition λ will be denoted $n_{\alpha}^k(\lambda)$.

The following lemma is the basis of the proof that the two crystals $B(\Lambda_0)$ and $B(\Lambda_0)^L$ are isomorphic.

Lemma 10.1.5. *Let λ be a node of $B(\Lambda_0)^L$. Let $\mathcal{L}$ be the k^{th} ladder of λ and $\mathcal{K}$ be the k^{th} ladder of $\mathcal{R}\lambda$. Let i be the residue of the ladders $\mathcal{L}$ and $\mathcal{K}$. Then the number of ladder-normal i -boxes on $\mathcal{L}$ is the same as the number of normal i -boxes on $\mathcal{K}$. Similarly, the number of ladder-conormal i -boxes on $\mathcal{L}$ is the same as the number of conormal i -boxes on $\mathcal{K}$. In particular, the ladder-good (ladder-cogood) i -box of λ lies on the same ladder as the good (cogood) i -box of $\mathcal{R}\lambda$.*

Proof. Suppose we are looking at the boxes on $\mathcal{L}$. Then note that:

- $n_b^k(\lambda) + \min(n_c^k(\lambda), n_d^k(\lambda)) = n_b^k(\mathcal{R}\lambda).$
- $n_e^k(\lambda) + \min(n_c^k(\lambda), n_d^k(\lambda)) = n_e^k(\mathcal{R}\lambda).$

This follows from the fact that a box of type (c) and a box of type (d) on $\mathcal{L}$ will combine under regularization to form a box of type (b) and a box of type (e) on $\mathcal{K}$.

Similarly,

- $n_g^k(\lambda) + \min(n_h^k(\lambda), n_i^k(\lambda)) = n_g^k(\mathcal{R}\lambda).$
- $n_j^k(\lambda) + \min(n_h^k(\lambda), n_i^k(\lambda)) = n_j^k(\mathcal{R}\lambda).$

Lemma 9.1.1 implies that there is no cancelation in the ladder i -signature on $\mathcal{L}$ (i.e. no type (e) node above a type (j) node). Similarly, Lemma 9.1.2 implies that there is no cancelation in the i -signature on $\mathcal{K}$ (i.e. no type (e) node below a type (j) node).

We also note that a box of type (b) on the k^{th} ladder of a partition implies that there is an addable i -box on the $(k + \ell)^{th}$ ladder, and that a box of type (g) on the k^{th} ladder of a partition implies that there is a removable i -box on the $(k - \ell)^{th}$ ladder.

$\mathcal{K}$ will contribute $\underbrace{+\cdots+}_{n_j^k(\lambda)} \underbrace{-\cdots-}_{n_e^k(\lambda)}$ to the ladder i -signature of λ . $\mathcal{L}$ will contribute $\underbrace{+\cdots+}_{n_j^k(\lambda)} \underbrace{+\cdots+}_{\min(n_h^k(\lambda), n_i^k(\lambda))} \underbrace{-\cdots-}_{n_e^k(\lambda)} \underbrace{-\cdots-}_{\min(n_c^k(\lambda), n_d^k(\lambda))}$ to the i -signature of $\mathcal{R}\lambda$. Since $\min(n_c^k(\lambda), n_d^k(\lambda)) = \min(n_h^{k+\ell}(\lambda), n_i^{k+\ell}(\lambda))$, the $\min(n_c^k(\lambda), n_d^k(\lambda))$ $-$'s will cancel with the $\min(n_h^{k+\ell}(\lambda), n_i^{k+\ell}(\lambda))$ $+$'s in the $k + \ell^{th}$ ladder, leaving just the $n_e^k(\lambda)$ $-$'s in the k^{th} ladder. Similarly, only the $n_j^k(\lambda)$ $+$'s will count in the i -signature of $\mathcal{R}\lambda$. Hence the number of normal and conormal boxes on any ladder is unchanged by regularization.

Since there are the same number of (co)normal boxes on each ladder of $\mathcal{R}\lambda$ as there are ladder-(co)normal boxes in each ladder of λ , the (co)good box of $\mathcal{R}\lambda$ must be on the same ladder as the ladder (co)good box of λ . $\square$

Corollary 10.1.6. *Let λ be a node in $B(\Lambda_0)^L$. Then $\widehat{\varphi}_i(\lambda) = \varphi_i(\mathcal{R}\lambda)$ and $\widehat{\varepsilon}_i(\lambda) = \varepsilon_i(\mathcal{R}\lambda)$.*

Theorem 10.1.7. *Regularization commutes with the crystal operators. In other words:*

- (1) $\mathcal{R} \circ \widehat{f}_i = \tilde{f}_i \circ \mathcal{R}$,
- (2) $\mathcal{R} \circ \widehat{e}_i = \tilde{e}_i \circ \mathcal{R}$.

Proof. (1) will follow from Lemma 10.1.5, since applying an $\widehat{f}_i$ to a partition λ will place an i -box in the same ladder of λ as $\tilde{f}_i$ places in $\mathcal{R}\lambda$. (2) follows similarly. $\square$

Corollary 10.1.8. *The crystal $B(\Lambda_0)$ is isomorphic to $B(\Lambda_0)^L$.*

Proof. The map $\mathcal{R} : B(\Lambda_0)^L \rightarrow B(\Lambda_0)$ gives the isomorphism. The map $\mathcal{S} : B(\Lambda_0) \rightarrow B(\Lambda_0)^L$ described in Section 7 is the inverse of $\mathcal{R}$. The other crystal isomorphism axioms are routine to check. $\square$

Example 10.1.9. Let $\lambda = (2, 1, 1, 1)$ and $\ell = 3$. Then $\mathcal{R}\lambda = (2, 2, 1)$. Also $\widehat{f}_2\lambda = (2, 1, 1, 1, 1)$ and $\tilde{f}_2(2, 2, 1) = (3, 2, 1)$. But $\mathcal{R}(2, 1, 1, 1, 1) = (3, 2, 1)$.

$$(2, 1, 1, 1) \xrightarrow{\widehat{f}_2} (2, 1, 1, 1, 1)$$

$$\mathcal{R} \downarrow \qquad \qquad \downarrow \mathcal{R}$$

$$(2, 2, 1) \xrightarrow{\tilde{f}_2} (3, 2, 1)$$

11. GENERALIZING CRYSTAL THEOREMS

We are now able to prove Theorems 4.3.5 and 4.3.6.

11.1. Crystal rules for weak ℓ -partitions. We first recall the following result due to James, which says that $D^{\mathcal{R}\lambda}$ occurs with multiplicity one in S^λ .

Theorem 11.1.1 (James [9]). *Let λ be any partition. Then $d_{\lambda, \mathcal{R}\lambda} = 1$.*

We can now prove our generalizations of Theorem 4.3.1 and Theorem 4.3.2.

THEOREM 4.3.5. *Suppose that λ is a weak ℓ -partition and $0 \leq i < \ell$. Then*

- (1) $\tilde{f}_i^\varphi \lambda$ is a weak ℓ -partition,
- (2) $\tilde{e}_i^\varphi \lambda$ is a weak ℓ -partition.

Proof. Let λ be a weak ℓ -partition. Then $D^\lambda = S^\nu$ for some $(\ell, 0)$ -JM partition ν with $\mathcal{R}\nu = \lambda$ (see Proposition 6.1.1). From Theorem 8.4.1 we know that $\nu \in B(\Lambda_0)^L$. Corollary 10.1.8 implies that $\widehat{\varphi}_i(\nu) = \varphi$. By Theorem 8.3.1, $\widehat{f}_i^\varphi \nu$ is another

($\ell, 0$)-JM partition. By Theorem 10.1.7 we know that $\mathcal{R}\widehat{f}_i^\varphi\nu = \widetilde{f}_i^\varphi\lambda$. Theorem 11.1.1 then implies that $D^{\widetilde{f}_i^\varphi\lambda} = S^{\widehat{f}_i^\varphi\nu}$, since $S^{\widehat{f}_i^\varphi\nu}$ is irreducible by Theorem 1.1.10. Hence $\widetilde{f}_i^\varphi\lambda$ is a weak ℓ -partition. (2) follows similarly. $\square$

THEOREM 4.3.6. *Suppose that λ is a weak ℓ -partition. Then*

- (1) $\widetilde{f}_i^k\lambda$ is not a weak ℓ -partition for $0 < k < \varphi - 1$,
- (2) $\widetilde{e}_i^k\lambda$ is not a weak ℓ -partition for $1 < k < \varepsilon$.

Proof. Let λ be a weak ℓ -partition. We must show that there does not exist an ($\ell, 0$)-JM partition μ in the regularization class of $\widetilde{f}_i^k\lambda$. There exists an ($\ell, 0$)-JM partition ν in $B(\Lambda_0)^L$ so that $D^\lambda = S^\nu$. By Theorem 8.3.2, $\widehat{f}_i^k\nu$ is not an ($\ell, 0$)-JM partition. But by Theorem 8.4.1 we know all ($\ell, 0$)-JM partitions occur in $B(\Lambda_0)^L$. Also, only one element of $\mathcal{RC}(\widetilde{f}_i^k\lambda)$ occurs in $B(\Lambda_0)^L$ and we know this is $\widehat{f}_i^k\nu$. Therefore no such μ can exist, so $\widetilde{f}_i^k\lambda$ is not a weak ℓ -partition. (2) follows similarly. $\square$

12. REPRESENTATION THEORETIC PROOF OF THEOREM 4.3.1

This proof relies heavily on the work of Grojnowski, Kleshchev et. al. We recall some notation from [7] but repeat very few definitions below.

12.1. Definitions and preliminaries. In the category Rep_n of finite dimensional representations of the finite Hecke algebra $H_n(q)$, we define the Grothendieck group $K(Rep_n)$ to be the abelian group generated by isomorphism classes of finite dimensional representations, with relations $[\mathcal{M}_1] + [\mathcal{M}_3] = [\mathcal{M}_2]$ if there exists an exact sequence $0 \rightarrow \mathcal{M}_1 \rightarrow \mathcal{M}_2 \rightarrow \mathcal{M}_3 \rightarrow 0$. This is a finitely generated group with generators corresponding to the irreducible representations of $H_n(q)$. The equivalence class corresponding to the module $\mathcal{M}$ is denoted $[\mathcal{M}]$.

Just as S_n can be viewed as the subgroup of S_{n+1} of permutations which fix $n+1$, $H_n(q)$ can be viewed as a subalgebra of $H_{n+1}(q)$ (the generators $T_1, T_2, \dots, T_{n-1}$ generate a subalgebra isomorphic to $H_n(q)$). Let $\mathcal{M}$ be a finite dimensional representation of $H_{n+1}(q)$. Then it makes sense to view $\mathcal{M}$ as a representation of $H_n(q)$. This module is called the *restriction of $\mathcal{M}$ to $H_n(q)$* , and is denoted $Res_{H_n(q)}^{H_{n+1}(q)}\mathcal{M}$. Similarly, we can define the induced representation of $\mathcal{M}$ by $Ind_{H_n(q)}^{H_{n+1}(q)}\mathcal{M} = H_{n+1}(q) \otimes_{H_n(q)} \mathcal{M}$.

To shorten notation, $Res_{H_b(q)}^{H_a(q)}$ will be written Res_b^a , and $Ind_{H_b(q)}^{H_a(q)}$ will be written as Ind_b^a .

If λ and μ are partitions, it is said that μ *covers* λ , (written $\mu \succ \lambda$) if the Young diagram of λ is contained in the Young diagram of μ and $|\mu| = |\lambda| + 1$.

Proposition 12.1.1. *Let λ be a partition of n and S^λ be the Specht module corresponding to λ . Then*

$$[Ind_n^{n+1} S^\lambda] = \sum_{\mu \succ \lambda} [S^\mu]$$

We consider functors $\tilde{e}_i : Rep_n \rightarrow Rep_{n-1}$ and $\tilde{f}_i : Rep_n \rightarrow Rep_{n+1}$ which commute with the crystal action on partitions in the following sense (see [7] for definitions and details).

Theorem 12.1.2. *Let λ be an ℓ -regular partition. Then*

- (1) $\tilde{e}_i D^\lambda = D^{\tilde{e}_i \lambda}$
- (2) $\tilde{f}_i D^\lambda = D^{\tilde{f}_i \lambda}$

We now consider the functors $f_i : Rep_n \rightarrow Rep_{n+1}$ and $e_i : Rep_n \rightarrow Rep_{n-1}$ which refine induction and restriction (for a definition of these functors, especially in the more general setting of cyclotomic Hecke algebras, see [7]). For a representation $\mathcal{M} \in Rep_n$ let $\varepsilon_i(\mathcal{M}) = \max\{k : e_i^k \mathcal{M} \neq 0\}$ and $\varphi_i(\mathcal{M}) = \max\{k : f_i^k \mathcal{M} \neq 0\}$. Grojnowski concludes the following theorem.

Theorem 12.1.3. *Let $\mathcal{M}$ be a finite-dimensional representation of $H_n(q)$. Let $\varphi = \varphi_i(\mathcal{M})$ and $\varepsilon = \varepsilon_i(\mathcal{M})$.*

- (1) $Ind_n^{n+1} \mathcal{M} = \bigoplus_i f_i \mathcal{M}; \quad Res_n^{n+1} \mathcal{M} = \bigoplus_i e_i \mathcal{M};$
- (2) $[f_i^\varphi \mathcal{M}] = \varphi! [\tilde{f}_i^\varphi \mathcal{M}]; \quad [e_i^\varepsilon \mathcal{M}] = \varepsilon! [\tilde{e}_i^\varepsilon \mathcal{M}].$

For a module D^μ the *central character* $\chi(D^\mu)$ can be identified with the multiset of residues of the partition μ .

Theorem 12.1.4. *All composition factors of the Specht module S^λ have the same central character.*

Theorem 12.1.5. $\chi(f_i(D^\lambda)) = \chi(D^\lambda) \cup \{i\}; \quad \chi(e_i(D^\lambda)) = \chi(D^\lambda) \setminus \{i\}.$

We are now ready to present a representation theoretic proof of Theorem 4.3.5, which states that if λ is a weak ℓ -partition lying at one end of an i -string in the crystal $B(\Lambda_0)$, then the opposite end of the i -string through λ is also a weak ℓ -partition.

12.2. A representation theoretic proof of Theorem 4.3.5.

THEOREM 4.3.5. *Suppose that λ is a weak ℓ -partition and $0 \leq i < \ell$. Then*

- (1) $\tilde{f}_i^\varphi \lambda$ is a weak ℓ -partition,
- (2) $\tilde{e}_i^\varepsilon \lambda$ is a weak ℓ -partition.

Alternate Proof of Theorem 4.3.5. Suppose λ is a weak ℓ -partition and $|\lambda| = n$. Recall that λ is a weak ℓ -partition if and only if there is a μ so that $S^\mu = D^\lambda$. Let F denote the number of addable i -boxes of μ and let ν denote the partition corresponding to μ plus all addable i -boxes.

First, we induce S^μ from $H_n(q)$ to $H_{n+F}(q)$. Applying Proposition 12.1.1 F times yields

$$[Ind_n^{n+F} S^\mu] = \sum_{\eta_F \succ \eta_{F-1} \succ \cdots \succ \eta_1 \succ \mu} [S^{\eta_F}].$$

Projecting to the direct summand which has the same central character as D^ν (i.e. the central character of μ with F more i 's), we get that $[S^\nu]$ occurs in this sum with coefficient $F!$ (add the i -boxes in any order). Everything else in this sum has a different central character.

We next apply (1) from Theorem 12.1.3 F times to obtain

$$[Ind_n^{n+F} D^\lambda] = \bigoplus_{i_1, \dots, i_F} [f_{i_1} \cdots f_{i_F} D^\lambda].$$

Projecting again to the direct summand with the same central character as ν , in $K(Rep)$ we get exactly $[(f_i)^F D^\lambda]$. Since λ is a weak ℓ -partition, $S^\mu = D^\lambda$, so $Ind_n^{n+F} S^\mu = Ind_n^{n+F} D^\lambda$ and we have shown that $F! [S^\nu] = [(f_i)^F D^\lambda]$.

Since $S^\mu = D^\lambda$ and $(f_i)^{\varphi+1} D^\lambda = 0$, we know that $F \leq \varphi$ ($Ind_n^{n+\varphi+1} D^\lambda$ has no direct summand with the correct central character by Theorem 12.1.5). Similarly, since $Ind_n^{n+F+1} S^\mu$ has no composition factors with central character $\chi(D^\lambda)$ union with $F+1$ many i 's, we know that $F \geq \varphi$. Hence $F = \varphi$.

By part 2 of Theorem 12.1.3, $[(f_i)^\varphi D^\lambda] = \varphi! [\tilde{f}_i^\varphi D^\lambda]$. Then by Theorem 12.1.2, $[S^\nu] = [D^{\tilde{f}_i^\varphi \lambda}]$.

Therefore $S^\nu = D\tilde{f}_i^\varphi \lambda$. Hence $\tilde{f}_i^\varphi \lambda$ is a weak ℓ -partition.

The proof that $\tilde{e}_i^{\varepsilon_i(\lambda)} \lambda$ is a weak ℓ -partition follows similarly, with the roles of induction and restriction changed in Proposition 12.1.1, and the roles of e_i and f_i changed in Theorem 12.1.3.

□

13. THE MULLINEUX MAP

The operation on the category of $H_n(q)$ modules of tensoring with the sign module is a functor which takes irreducible modules to irreducible modules. For instance, when q is not a root of unity, then $S^\lambda \otimes \text{sign} \cong S^{\lambda'}$. When λ is an ℓ -regular partition, and D^λ denotes the irreducible module corresponding to λ then $D^\lambda \otimes \text{sign}$ is some irreducible module $D^{m(\lambda)}$. This describes a map m on ℓ -regular partitions called the *Mullineux map*. Recent results of Fayers [5] settle a conjecture of Lyle [16] which effectively computes the Mullineux map in certain cases by means of regularization and transposition. This section will highlight the interpretation of Fayers result in terms of the ladder crystal. It should be noted that Ford and Kleshchev gave a recursive construction for computing the Mullineux map [6].

13.1. Statement of Fayers Results. Since the Mullineux map is the modular analog of transposition, a natural attempt to compute $m(\lambda)$ for an ℓ -regular partition λ would be to transpose λ and then regularize. If a partition is not ℓ -regular, we could similarly guess that $m(\mathcal{R}\lambda)$ was the composition of transposition and regularization. This is not always the case. However, a conjecture of Lyle [16], which was proven recently by Fayers [5] gives a precise classification for when this holds. The definition below was taken from Fayers [5].

Definition 13.1.1. An L -partition is a partition which has no box (i, j) in the diagram of λ such that $\ell \mid h_{i,j}^\lambda$ and either $\text{arm}(i, j) < (\ell - 1) * \text{leg}(i, j)$ or $\text{leg}(i, j) < (\ell - 1) * \text{arm}(i, j)$.

Theorem 13.1.2. [Fayers [5]] *A partition is an L -partition if and only if $m(\mathcal{R}\lambda) = \mathcal{R}(\lambda')$.*

13.2. All L-partitions occur as nodes in the ladder crystal. Now we prove that all L-partitions are nodes in the crystal $B(\Lambda_0)^L$. To do this, we will use an equivalent condition of L-partitions given by Lyle in [16].

Lemma 13.2.1. *A partition λ is an L-partition if and only if there does not exist a box (i, j) in the diagram of λ such that $\ell \mid h_{(i,j)}^\lambda$, and $\frac{h_{(i,j)}^\lambda}{\ell} \leq \min\{\text{arm}(i, j), \text{leg}(i, j)\}$.*

Proof. Suppose that λ is not an L-partition. Let (i, j) be such that $\ell \mid h_{(i,j)}^\lambda$ and suppose that $\text{arm}(i, j) < (\ell - 1) * \text{leg}(i, j)$ (the case where $\text{leg}(i, j) < (\ell - 1) * \text{arm}(i, j)$ is similar). Then $h_{i,j}^\lambda = \text{arm}(i, j) + \text{leg}(i, j) + 1 < \ell * \text{leg}(i, j) + 1$. So $\frac{h_{(i,j)}^\lambda}{\ell} < \text{leg}(i, j) + \frac{1}{\ell}$, or $\frac{h_{(i,j)}^\lambda}{\ell} \leq \text{leg}(i, j)$. Therefore $\frac{h_{(i,j)}^\lambda}{\ell} \leq \min\{\text{arm}(i, j), \text{leg}(i, j)\}$.

Now suppose there exists a box (i, j) so that $\ell \mid h_{(i,j)}^\lambda$, and

$$\frac{h_{(i,j)}^\lambda}{\ell} \leq \min\{\text{arm}(i, j), \text{leg}(i, j)\}.$$

Let $k = \min\{\text{arm}(i, j), \text{leg}(i, j)\}$. If $k = \text{arm}(i, j)$, then $h_{i,j}^\lambda \leq \ell * \text{arm}(i, j)$, so $\text{leg}(i, j) = h_{i,j}^\lambda - \text{arm}(i, j) - 1 \leq (\ell - 1) * \text{arm}(i, j) - 1$. Therefore $\text{leg}(i, j) < (\ell - 1) * \text{arm}(i, j)$. The $k = \text{leg}(i, j)$ case follows similarly. $\square$

Theorem 13.2.2. *All L-partitions occur as nodes of the crystal $B(\Lambda_0)^L$.*

Proof. Suppose that λ is not a node of the crystal $B(\Lambda_0)^L$. Then there exists an unlocked box somewhere in the diagram of λ , say at position (a, b) . Let c be the smallest integer so that (c, b) is unlocked (so that either $(c - 1, b)$ is locked or $c = 1$). Then the box (c, λ_c) is also unlocked (if it were locked then (c, b) would be a type II locked box). If there is an unlocked box above (c, λ_c) then we will apply this procedure again (move to the highest unlocked box in column λ_c and then the rightmost box in that row) until we can assume that (c, λ_c) is unlocked and sits below a locked box (or possibly $c = 1$). Then (c, λ_c) must be unlocked from violating the type I lock condition, so that there exists a position (i, j) in the same ladder as (c, λ_c) such that (i, j) is not in λ but $(i - 1, j)$ is in λ . Then the hook length $h_{(c,j)}^\lambda$ is divisible by ℓ . In fact, $h_{(c,j)}^\lambda = \ell * \text{arm}(c, j)$, since (i, j) is in the same ladder as (c, λ_c) and (i, j) is the first non-occupied position in column j in λ . But then $\frac{h_{(c,j)}^\lambda}{\ell} = \text{arm}(c, j)$ and $\text{leg}(c, j) = (\ell - 1) * \text{arm}(c, j) - 1 \geq \text{arm}(c, j)$ since $\ell > 2$. Hence $\frac{h_{(c,j)}^\lambda}{\ell} \leq \min(\text{arm}(c, j), \text{leg}(c, j))$, which means λ is not an L-partition. $\square$

It was pointed out to the author by M. Fayers that a classification of the nodes of $B(\Lambda_0)^L$ can be described in terms of hook lengths and arm lengths. We now include this classification.

Theorem 13.2.3. *A partition λ belongs to the crystal $B(\Lambda_0)^L$ if and only if there does not exist a box $(i, j) \in \lambda$ such that $h_{(i,j)}^\lambda = \ell * \text{arm}(i, j)$.*

Proof. If λ has a box (i, j) with hook length $\ell * \text{arm}(i, j)$ then the box (i, λ_i) will be unlocked, as it is on the same ladder as the empty position directly below the last box in column j . Hence λ is not in $B(\Lambda_0)^L$.

If λ does not belong to the crystal $B(\Lambda_0)^L$ then there exists an unlocked box (a, b) in the diagram of λ . Similar to the proof of Theorem 13.2.2, this implies that there exists an unlocked box (i, λ_i) which is either directly below a locked box or $i = 1$. This box is then unlocked because it violates a type I lock rule, which means that in the same ladder as (i, λ_i) there is an empty position (n, j) directly below a box $(n - 1, j)$. But then the box (i, j) will satisfy $h_{(i,j)}^\lambda = \ell * \text{arm}(i, j)$. $\square$

Finally we note that as a corollary we obtain a second proof that all $(\ell, 0)$ -JM partitions are nodes of the crystal $B(\Lambda_0)^L$.

Corollary 13.2.4. *All $(\ell, 0)$ -JM partitions (in fact, all (ℓ, p) -JM partitions) are nodes of the crystal $B(\Lambda_0)^L$.*

Proof. The two composition factors $D^{\mathcal{R}(\lambda')}$ and $D^{m(\mathcal{R}\lambda)}$ must occur in the Specht module $S^{\lambda'}$. If λ is a JM partition then so is λ' , so $S^{\lambda'}$ is irreducible, i.e. $S^{\lambda'} \cong D^{\mathcal{R}(\lambda')} = D^{m(\mathcal{R}\lambda)}$. By Theorem 13.1.2, λ is an L-partition. Hence all JM-partitions are L-partitions. $\square$

REFERENCES

- [1] C. Berg and M. Vazirani, $(\ell, 0)$ -Carter partitions, a generating function, and their crystal theoretic interpretation, *ArXiv Mathematics e-prints*, math.CO/0712.2075.
- [2] J.P. Cossey, M. Ondrus, and C.R. Vinroot, Constructing all irreducible Specht modules in a block of the symmetric group, *ArXiv Mathematics e-prints*, math/0605654
- [3] R. Dipper and G. James, *Representations of Hecke algebras of general linear groups*, Proc. LMS (3), **52** (1986), 20-52
- [4] M. Fayers, Irreducible Specht modules for Hecke Algebras of Type A, *Adv. Math.* **193** (2005) 438-452

- [5] M. Fayers, Regularisation and the Mullineux map, *ArXiv Mathematics e-prints*, math.CO/0808.1679
- [6] B. Ford and A. Kleshchev, A proof of the Mullineux conjecture, *Math. Z.* **226** (1997), 267-308.
- [7] I. Grojnowski, Affine $\widehat{\mathfrak{sl}}_p$ controls the representation theory of the symmetric group and related Hecke algebras, *ArXiv Mathematics e-prints*, math/9907129
- [8] G.D. James, The decomposition matrices of $GL_n(q)$ for $n \leq 10$, *Proc. Lond. Math. Soc.* (3), **60** (1990), 225-264.
- [9] G.D. James, On the Decomposition Matrices of the Symmetric Groups II, *J. of Algebra* **43** (1976), 45-54.
- [10] G.D. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Encyclopedia of Mathematics, **16**, 1981.
- [11] G.D. James and A. Mathas, A q -analogue of the Jantzen-Schaper theorem, *Proc. Lond. Math. Soc.*, **74** (1997), 241-274.
- [12] M. Kashiwara, On crystal bases, in Representations of groups (Banff 1994), *CMS Conf. Proc.* **16** (1995), 155-197.
- [13] A. Kleshchev, *Linear and Projective Representations of Symmetric Groups*, Cambridge Tracts in Mathematics **163**.
- [14] A. Lascoux, B. Leclerc, and J.-Y. Thibon, Hecke algebras at roots of unity and crystal bases of quantum affine algebras, *Comm. Math. Phys.* **181** (1996), 205-263.
- [15] S. Lyle, Some q -analogues of the Carter-Payne Theorem, *J. Reine Angew. Math.*, to appear
- [16] S. Lyle, Some topics in the representation theory of the symmetric and general linear groups, Ph.D. thesis, University of London, 2003.
- [17] A. Mathas, *Iwahori-Hecke algebras and Schur algebras of the symmetric group*, University Lecture Series, **15**, AMS, 1999
- [18] K.C. Misra and T. Miwa, Crystal base for the basic representation of $U_q(\mathfrak{sl}_n)$, *Commun. Math. Phys.* **134** (1990), 79-88.
- [19] B. E. Sagan, *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, Springer GTM, 2001.

CHRIS BERG, DEPARTMENT OF MATHEMATICS, UC DAVIS, USA, 95616

E-mail address: `berg@math.ucdavis.edu`